

SOME NEW RESULTS ON CIRCLE GRAPHS

Guillermo Durán*

Abstract

The intersection graph of a set of chords on a circle is called a circle graph. This class of graphs admits some interesting subclasses, such as: Helly circle graphs, clique-Helly circle graphs, unit circle graphs and proper circular-arc graphs. In this paper, we prove some inclusion relations among these subclasses. A necessary condition for a graph to be a Helly circle graph is shown and we conjecture that this condition is sufficient too. All possible intersections among these subclasses are analyzed. The number of regions generated is 10. We show a minimal example belonging to each one of them. Finally, some properties about minimal forbidden subgraphs for circle graphs are proved.

1 Introduction

Consider a finite family of non-empty sets. The intersection graph of this family is obtained by representing each set by a vertex, two vertices being connected by an edge if and only if the corresponding sets intersect. Intersection graphs have received much attention in the study of algorithmic graph theory and their applications [8]. Well known special classes of intersection graphs include interval graphs, chordal graphs, permutation graphs, circle graphs, circular-arc graphs, and so on.

A graph G is called a circle graph (CG) if there exists a set of chords L (a model) on a circle and a one-to-one correspondence between vertices of G and chords of L such that two distinct vertices are adjacent if and only if their corresponding chords intersect in the interior of the circle. That is, a circle graph is the intersection graph of a set of chords on a circle. Figure 1 shows a circle graph G and a model L for it.

*Partially supported by FONDECYT Grant 1030498 and Millennium Science Nucleus “Complex Engineering Systems”, Chile and UBACyT Grant X127, Argentina.

Keywords: circle graphs, clique-Helly circle graphs, Helly circle graphs, proper circular-arc graphs, unit circle graphs.




Figure 1

Circle graphs were introduced in [6], where an application to solve a problem about queues and stacks proposed by Knuth [9] is shown. Polynomial time algorithms for recognizing graphs in this class appear in [1, 7, 10, 11].

A family of subsets S satisfies the Helly property when every subfamily of it consisting of pairwise intersecting subsets has a common element. A graph is clique-Helly when its cliques (maximal complete subgraphs) satisfy the Helly property. A circular-arc graph is the intersection graph of a family of arcs on a circle (we assume that all the arcs are open). A graph is diamond-free if it does not contain a diamond (Figure 2) as an induced subgraph.




Figure 2

A local complementation of a graph G at vertex v consists in replacing $G_{N[v]}$ by its complement graph (where $G_{N[v]}$ is the induced subgraph by the neighbours of v). Graphs G_1 and G_2 are locally equivalent if there is a finite sequence of local complementations transforming G_1 into G_2 .

Let \overline{G} be the complement of a graph G and G^* , a graph G adding it an isolated vertex. Let C_j be an induced cycle with j vertices.

The proposal of this paper is to analyze different subclasses of circle graphs and find some properties about minimal forbidden subgraphs for this class.

In Section 2, some inclusion relations among the subclasses of circle graphs

that we will define are proved. A necessary condition for a graph to be a Helly circle graph is also shown and we conjecture that this condition is sufficient too.

In Section 3, all possible intersections among these subclasses of circle graphs are analyzed, as it was done in [5] for circular-arc graphs and their subclasses: clique-Helly circular-arc graphs, Helly circular-arc graphs, proper circular-arc graphs and unit circular-arc graphs. The number of regions generated is 10. We show a minimal example belonging to each one of them. The minimality of the examples implies that any proper induced subgraph of them belongs to some other region.

In Section 4, a characterization of proper circular-arc graphs by forbidden subgraphs due to Tucker [14] is used to show some properties about minimal forbidden subgraphs for circle graphs.

2 Subclasses of circle graphs

We may define some interesting subclasses of circle graphs:

- (i) Helly circle graphs: a graph G is a Helly circle (HC) graph if there is a model L for G such that the set of chords of L satisfies the Helly property. It is not known a characterization of this subclass.
- (ii) Clique-Helly circle graphs: a graph G is a clique-Helly circle (CHC) graph if G is a circle graph and a clique-Helly graph. Szwarcfiter [12] described a characterization of clique-Helly graphs leading to a polynomial time algorithm for recognizing them. This algorithm together with an efficient method for circle graphs results in a polynomial time algorithm for recognizing CHC graphs.
- (iii) Unit circle graphs: a graph G is a unit circle (UC) graph if there is a model L for G such that all the chords are of the same length. We will see that this subclass is equivalent to unit circular-arc (UCA) graphs, circular-arc graphs which have a representation where all the arcs are of the same length. These graphs can be also recognized in polynomial time [4].
- (iv) Proper circular-arc graphs: a graph G is a proper circular-arc (PCA) graph if there is a circular-arc representation of G such that no arc is

properly contained in any other. It can be easily proved that proper circular-arc graphs are a subclass of circle graphs. The representation in arcs can be trivially transformed in the model in chords, using that no pair of arcs together covers the entire circle (a property of proper circular-arc graphs shown in [8]). Tucker [13] described a characterization and an efficient algorithm for recognizing *PCA* graphs.

First, we prove a necessary condition so that a graph is a Helly circle graph.

Theorem 2.1 *Let G be a Helly circle graph. Then, G is a diamond free circle graph.*

Proof: Let G be a Helly circle graph and suppose that G contains a diamond D (Figure 2) as an induced subgraph. Given $V(D) = \{v_1, v_2, v_3, v_4\}$ and let L_1, L_2, L_3 and L_4 be four different chords which verify Helly property, corresponding to the vertices v_1, v_2, v_3 and v_4 , respectively. As the chords verify Helly property, L_1, L_2 and L_3 must have a common point P because v_1, v_2 and v_3 define a clique of D . Analogously, L_2, L_3 and L_4 must have a common point Q because v_2, v_3 and v_4 define another clique of D . If $P \neq Q$, then $L_2 = L_3$, a contradiction. If $P = Q$, then D is the clique with 4 vertices, a contradiction again.

□

We conjecture that the converse of this theorem holds too. So, the diamond would be the only minimal obstruction to a circle graph to be a Helly circle graph. If this conjecture were true, we would have a polynomial recognition of the *HC* subclass, because we can check in polynomial time whether a graph has a diamond as an induced subgraph.

Now, we prove that *HC* is a subclass of *CHC*.

Theorem 2.2 *Let G be a Helly circle graph. Then, G is a clique-Helly circle graph.*

Proof: It was proved in [3] that diamond-free graphs are a subclass of clique-Helly graphs. And by Theorem 2.1, Helly circle graphs are included in diamond-free circle graphs. So, this theorem holds.

□

It is interesting to remark that the analogous result for circular-arc graphs is not true. Helly circular-arc graphs are not necessarily clique-Helly [5].

Finally, it can be easily proved that unit circle graphs are equivalent to unit circular-arc graphs.

Theorem 2.3 *A graph G is a unit circle graph if and only if G is a unit circular-arc graph.*

Proof: Let us see the equivalence. The model in chords of an UC graph G can be transformed into a circular-arc representation of G in arcs of equal length using the same endpoints and joining them such that the arc has length at most π (assuming the radius $r = 1$). The converse also holds. Let G be an UCA graph. It is easy to see that we have a representation in unit arcs of G without common endpoints. We may suppose, without loss of generality, that each arc has length at most π , assuming again the radius $r = 1$ (otherwise, the graph is complete and, in consequence, it is a unit circle graph). Then, if we join the endpoints of each arc we have a circle model of G in chords of the same length. \square

A trivial consequence of Theorem 2.3 is the following inclusion.

Corollary 2.1 *Let G be a unit circle graph. Then, G is a proper circular-arc graph.*

3 Minimal examples

In Figure 4 we can see minimal examples of graphs belonging to the possible intersections defined by the relations among these classes. The examples can be checked with no difficulty by the reader. We present here only one proof of a minimal member belonging to the respective region.

First, we need some characterization theorems.

In order to characterize the $PCA - UCA$ region (which is equivalent to the $PCA - UC$ region), we need a definition due to Tucker [14], which was formalized in [5].

Let j and k be two positive integers with $j > k$. The graph $CI(j, k)$ is the circular-arc graph corresponding to the circular-arc representation built in the following way: let ε be a positive real number, and $r=1$ the radius of the circle. Draw j arcs A_0, A_1, \dots, A_{j-1} of length $l_1 = 2\pi k/j + \varepsilon$ such that each A_i starts at $2\pi i/j$ and ends at $2\pi(i+k)/j + \varepsilon$ (i.e., $A_i = (2\pi i/j, 2\pi(i+k)/j + \varepsilon)$). Afterwards, draw j arcs B_0, B_1, \dots, B_{j-1} of length $l_2 = 2\pi k/j - \varepsilon$, such that

each B_i starts at $(2\pi i + \pi)/j$ and ends at $(2\pi(i + k) + \pi)/j - \varepsilon$ (i.e., $B_i = ((2\pi i + \pi)/j, (2\pi(i + k) + \pi)/j - \varepsilon)$). For example, the representation of Figure 3 generates the graph $CI(4, 1)$.




Figure 3

By construction, these graphs are proper circular-arc graphs [14].

Theorem 3.1 [14] *Let G be a proper circular-arc graph. Then G is a unit circular-arc graph if and only if G contains no $CI(j, k)$ as an induced subgraph, where j and k are relatively prime and $j > 2k$.*

It can be proved that this characterization of UCA graphs leads to a polynomial time algorithm for the recognition of this class [4].

A characterization of clique-Helly graphs is presented in [12]. We must define the concept of an extended triangle. Let G be a graph and T a triangle of G . The extended triangle of G , relative to T , is the subgraph of G induced by all the vertices which form a triangle with at least one edge of T . A vertex v is universal in a subgraph of G if v is adjacent to every other vertex of the subgraph.

Theorem 3.2 [12] *A graph G is a clique-Helly graph if and only if every of its extended triangles contains a universal vertex.*

Now, we can prove the following result.

Proposition 3.1 *Graph H (Figure 5) is a proper circular-arc graph and a clique-Helly circle graph but it is neither a unit circle graph nor a Helly circle graph.*




Figure 4




Figure 5

Proof:

- (i) Graph H is a proper circular arc graph but it is not a unit circle graph: by the definition of the graphs $CI(j, k)$, H is isomorphic to $CI(7, 2)$ and this graph belongs to the $PCA - UCA$ region.
- (ii) Graph H is a clique-Helly circle graph: by inspection, we can verify that every extended triangle of H contains a universal vertex. So, by Theorem 3.2, H is clique-Helly.
- (iii) Graph H is not a Helly circle graph: the subgraph induced by v_1, v_2, v_3 and v_4 is a diamond.

In order to verify the minimality of this example we prove that any proper induced subgraph H' of H is UCA . With this purpose, we show that H' does not contain $CI(j, k)$ as an induced subgraph, where j and k are relatively prime and $j > 2k$. It is enough to prove this fact for $CI(3, 1)$, $CI(4, 1)$, $CI(5, 1)$, $CI(6, 1)$ and $CI(5, 2)$ because they are the graphs of this family with at most thirteen vertices.

Since H has a maximum independent set of size three, so H' cannot contain $CI(4, 1)$, $CI(5, 1)$ and $CI(6, 1)$, which have maximum independent set of size 4, 5 and 6, respectively. Now, suppose that H' contains $CI(3, 1)$ as an induced subgraph. Observe that there is no stable set of size three induced by the vertices v_i 's (since the subgraph induced by all v_i 's is the complement of C_7). So by symmetry, we may assume that every stable set contains w_3 and we have to analyze only two cases: the stable set of size three in $CI(3, 1)$ is formed either by $\{v_1, w_3, w_5\}$ or by $\{w_1, w_3, w_5\}$.

In the first case (Figure 6), the vertices which are adjacent to v_1 and w_5 simultaneously and non adjacent to w_3 , are v_6, v_7 and w_6 ; adjacent to v_1 and w_3 and not to w_5 , are v_2, v_3 and w_2 ; and adjacent to w_3 and w_5 and not to v_1 , are v_4, v_5 and w_4 . But we cannot choose three of these vertices such that they induce a triangle. Then, H' cannot contain $CI(3, 1)$ as an induced subgraph.

In the second case (Figure 7), the vertex adjacent to w_1 and w_5 and non adjacent to w_3 must be v_7 ; the vertex adjacent to w_1, w_3 and v_7 and non adjacent to w_5 must be v_2 . But now, we don't have a vertex adjacent to v_2, v_7, w_3 and w_5 , and non adjacent to w_1 . So, H' cannot contain $CI(3, 1)$ as an induced subgraph.




Figure 6




Figure 7

Finally, suppose that H' contains $CI(5, 2)$ as an induced subgraph. H' must contain the structure of Figure 8 included in $CI(5, 2)$ (for more detail about this graph, see [5]):




Figure 8

By the symmetry of $CI(7, 2)$, we may assume, without loss of generality, that one of the K_5 is induced by v_1, v_2, v_3, w_1 and w_2 . But now, there is not any vertex in $CI(7, 2)$ adjacent to exactly four of these vertices. So, H' does not contain $CI(5, 2)$ as an induced subgraph.

□

4 Minimal forbidden subgraphs

It is not known a characterization of circle graphs by forbidden subgraphs. An interesting approach in that sense is a characterization of this class proved by Bouchet [2].

Theorem 4.1 [2] *A graph G is a circle graph if and only if no graph locally equivalent to G contains an induced subgraph isomorphic either to W_5 or W_7 or BW_3 (Figure 9).*




Figure 9

A trivial corollary of this theorem and the transitivity of the local equivalence is the following.

Corollary 4.1 *Let H be a (not) circle graph. If a graph G is locally equivalent to H , then G is (not) a circle graph.*

A characterization of proper circular-arc graphs by forbidden subgraphs due to Tucker [14] is used here to show some properties about minimal forbidden subgraphs for circle graphs. First, we need the definition of H_1, H_2, H_3, H_4 and H_5 , the graphs of Figure 10.

Theorem 4.2 [14] *A graph G is a proper circular-arc graph if and only if it contains as induced subgraphs none of these graphs: $\overline{H_1}, \overline{H_2}, \overline{H_3}, \overline{H_4}, \overline{H_5}, H_1^*, \{C_n^*\}_{n \geq 4}, \{\overline{C_{2j+1}^*}\}_{j \geq 1}$ and $\{\overline{C_{2j}^*}\}_{j \geq 3}$.*

We have to analyze which of the forbidden subgraphs of Theorem 4.2 are circle graphs and which are not.

Lemma 4.1 *Graphs $\overline{H_1}, \overline{H_3}, H_1^*, \{\overline{C_3^*}\}$ and $\{C_n^*\}_{n \geq 4}$ are circle graphs.*




Figure 10




Figure 11

Proof: It is easy to find circle models for $\overline{H_1}, H_1^*, \{\overline{C_3^*}\}$ and $\{C_n^*\}_{n \geq 4}$. Let us see the circle model for $\overline{H_3}$ (Figure 11)

So, the lemma holds. □

Lemma 4.2 *Graphs $\overline{H_2}, \overline{H_4}, \overline{H_5}, \{\overline{C_{2j+1}^*}\}_{j \geq 2}$ and $\{\overline{C_{2j}^*}\}_{j \geq 3}$ are not circle graphs.*

Proof:

- (i) Graph $\overline{H_2}$ (Figure 12) is not a circle graph. If we apply the operation of local complementation first at vertex v_4 and then at vertex v_1 , we obtain a graph isomorphic to BW_3 .




Figure 12

- (ii) Graph $\overline{H_4}$ (Figure 13) is not a circle graph. If we apply the operation of local complementation first at vertex v_6 and then at vertex v_3 , the subgraph induced by vertices v_1, v_2, v_4, v_5, v_6 and v_7 is isomorphic to W_5 .




Figure 13

- (iii) Graph $\overline{H_5}$ (Figure 14) is not a circle graph. If we apply the operation of local complementation successively at vertices v_5, v_3 and v_7 , we obtain a graph isomorphic to BW_3 again.




Figure 16

The following facts assert that one of the extreme points of L_{2k+1} is between a_{2k-1} and b_1 , and the other one is between b_{2k-1} and a_1 :

- Vertex v_{2k+1} forms a complete subgraph with $v_1, v_3, \dots, v_{2k-1}$.
- Vertex v_{2k} is adjacent to any odd vertex, except v_{2k-1} and v_{2k+1} .
- Vertex v_{2k-2} is adjacent to any odd vertex, except v_{2k-3} and v_{2k-1} .

We have to draw now even chords. Analogously, they verify the same property of odd chords. First, draw the chord L_2 corresponding to vertex v_2 . As v_2 is adjacent to any odd vertex except v_1 and v_3 , we may assume without loss of generality that its corresponding chord has one of the extremes between a_3 and a_5 , and the other one between a_{2j-1} and b_1 . Repeat this analysis for $L_4, L_6, \dots, L_{2j-2}$ and conclude that we cannot draw the chord L_{2j} .

□

Now, the following result about minimal forbidden subgraphs for circle graphs can be proved.

Theorem 4.3 *Let H be a minimal forbidden subgraph for circle graphs. Then either H contains properly as an induced subgraph some of the graphs of Lemma 4.1 or H is isomorphic to some of the graphs of Lemma 4.2.*

Proof: It is a direct consequence of Theorem 4.2, Lemmas 4.1 and 4.2, and the fact of proper circular-arc graphs are a subclass of circle graphs.

□

Acknowledgments. To the anonymous referees for their interesting observations, to Jayme Szwarcfiter for his comments and suggestions which improved this work, and to Flavia Bonomo for her ideas to prove one of the parts of Lemma 4.2 and her help during the preparation of the paper.

References

- [1] Bouchet, A., *Reducing prime graphs and recognizing circle graphs*, *Combinatorica* 7, 3 (1987), 243-254.
- [2] Bouchet, A., *Circle graph obstructions*, *Journal of Combinatorial Theory B* 60 (1994), 107-144.
- [3] Chong-Keang, L.; Yee-Hock, P., *On graphs without multicliqual edges*, *Journal of Graph Theory* 5 (1981), 443-451.
- [4] Durán, G.; Gravano, A.; Spinrad, J.; Tucker, A., *Polynomial time recognition of unit circular-arc graphs*, submitted.
- [5] Durán, G.; Lin, M., *On some subclasses of circular-arc graphs*, *Congressus Numerantium* 146 (2000), 201-212.
- [6] Even, S.; Itai, A., *Queues, stacks and graphs*, In *Theory of Machines and Computations*, A. Kohavi and A.Paz eds., Acad. Press, N. York (1971), 71-86.
- [7] Gabor, C.; Supowit, K.; Hsu, W., *Recognizing circle graphs in polynomial time*, *Journal of the ACM*, Vol. 36, 3 (1989), 435-473.
- [8] Golumbic, M., *Algorithm Graph Theory and Perfect Graphs* (Academic Press, New York, 1980).
- [9] Knuth, D., *The Art of Computer Programming*, Vol. 1, Addison-Wesley, (1969).
- [10] Naji, W., *Reconnaissance des graphes de cordes*, *Discrete Math.* 54 (1985), 329-337.
- [11] Spinrad, J., *Recognition of circle graphs*, *Journal of Algorithms* 16 (2) (1994), 264-282.

- [12] Szwarefiter, J., *Recognizing clique-Helly graphs*, Ars Combinatoria 45 (1997), 29-32.
- [13] Tucker, A., *Matrix characterizations of circular-arc graphs*, Pacific J. Math. 38 (1971), 535-545.
- [14] Tucker, A., *Structure theorems for some circular-arc graphs*, Discrete Mathematics 7 (1974), 167-195.

Departamento de Ingeniería Industrial
Facultad de Ciencias Físicas y Matemáticas
Universidad de Chile
Santiago, Chile
`gduran@dii.uchile.cl`

Departamento de Computación
Facultad de Ciencias Exactas y Naturales
Universidad de Buenos Aires
Buenos Aires, Argentina