

# On the Helly Number in $P_3$ -Convexity in Graphs

Moisés T. Carvalho      Mitre C. Dourado  
Jayme L. Szwarcfiter

## Abstract

A subset of vertices  $S \subseteq V(G)$  of a graph  $G$  is  $P_3$ -convex if all vertices with at least two neighbors on  $S$  lie on  $S$ . The *Helly number* is the least integer  $k$  such that any subfamily of  $k$ -intersecting  $P_3$ -convex sets of  $G$  contains a common vertex. In this work, we determine the Helly number for path graphs, cycles, and threshold graphs. We show that complete  $k$ -partite graphs satisfies the Helly property and we present bounds for the Helly number of a graph in the  $P_3$ -convexity.

## 1 Introduction

Given a finite set  $V$ , a family  $\mathcal{C}$  of subsets of  $V$  is a *convexity* in  $V$  if:

- (1) The empty set and  $V$  belong to  $\mathcal{C}$ .
- (2)  $\mathcal{C}$  is closed under intersection.

The subsets of  $V$  in  $\mathcal{C}$  are called *convex sets*.

---

*2000 AMS Subject Classification:* 05C12, 68R10.

*Key Words and Phrases:* Convexity,  $P_3$ -convexity, Helly number.

Support by CAPES and CNPq

In this work, we consider a convexity in graphs known as  $P_3$ -convexity [3]. Let  $G$  be a graph and  $S$  a set of vertices of  $V(G)$ ,  $S$  is  $P_3$ -convex if every vertex with at least two neighbors on  $S$  belongs to  $S$ . The *closed interval* between two vertices  $u$  and  $v$  is the set  $I_{P_3}[u, v]$  (or  $I[u, v]$ , if there is no ambiguity) that contains  $u$  and  $v$  and all vertices of paths of length two between the pair of vertices  $u, v$ , and the *open interval*  $I(u, v) = I[u, v] \setminus \{u, v\}$ . The set  $\langle S \rangle$  is the smallest convex set of  $G$  containing  $S$  and is called the *convex hull* of  $S$  in  $G$  [6].

The *core* of a family of sets  $\mathcal{C}$  is defined as  $core(\mathcal{C}) = E_1 \cap E_2 \cap \dots \cap E_m$  where  $E_i \subseteq V(G)$ , for  $i = 1, \dots, m$ . A family of sets is  $k$ -*intersecting* if each  $k$  of their sets has a non-empty core. A family  $\mathcal{C}$  is  $k$ -*Helly* when every  $k$ -intersecting subfamily of  $\mathcal{C}$  has a non-empty core. The *Helly number* of  $\mathcal{C}$  is the smallest integer  $p$  for which  $\mathcal{C}$  is  $p$ -Helly. When  $p = 2$ , we say that  $\mathcal{C}$  satisfies the *Helly property* [2]. The Helly property got its name thanks to the mathematician Eduard Helly [5]. We denote by  $\mathcal{M}_G$  the set with all convex sets in the graph  $G$  and denote by  $G[A]$  a subgraph of  $G$  induced by  $A$ , where  $A \subseteq V(G)$ .

In this paper we study the Helly number in  $P_3$ -convexity in graphs, denoted by  $h(G)$ .

The next Theorem by Berge and Duchet [1] and the Lemma by Berge, Duchet and Calder [4] will be very useful in our proofs.

**Theorem 1** (Berge and Duchet [1]). *A family of sets  $\mathcal{C}$  is  $k$ -Helly if and only if for every set  $A \subseteq V(\mathcal{C})$  with  $|A| = k + 1$ , the intersection of the sets  $S_j$ , with  $|S_j \cap A| \geq k$ , is non-empty.*

**Lemma 2** (Berge, Duchet and Calder [4]). *In any convexity, the Helly number is the least integer  $k$  such that every  $(k + 1)$ -element set  $S \subseteq V$  has the property  $\bigcap_{a \in S} \langle S \setminus \{a\} \rangle \neq \emptyset$ .*

In the graph shown in Figure 1, we have two different sets formed by black vertices. The first one is a convex set and the second one is not a convex set.



Figure 1: Example and counterexample of  $P_3$ -convex set.

## 2 The Helly number in $P_3$ -convexity

In this section, we present a theorem about universal vertices, and results to some graph classes. In all proofs, the trivial graph is not considered since its Helly number is equal to one. Also we only consider connected graphs.

**Lemma 3.** *Let  $G$  be a graph. If all non-empty convex sets  $S_i$  of  $\mathcal{M}_G$  are  $S_i = \{v_i\}$  (vertices of  $G$ ),  $S_i = \{u, v\}$  (induces an edge in  $G$ ) or  $S_i = V(G)$ , then  $h(G) = 2$ .*

*Proof.* If  $G$  is isomorphic to  $K_3$ , it is easy to see that  $h(G) = 2$ .

Assume  $G$  is not isomorphic to  $K_3$ . Suppose for contradiction that  $h(G) \geq 3$ . Then there is  $A \subseteq V(G)$ ,  $A = \{a_1, \dots, a_k\}$ ,  $|A| \geq 3$  such that  $\bigcap_{a_i \in A} \langle A \setminus \{a_i\} \rangle = \emptyset$ . We can assume that  $S_j = \langle A \setminus \{a_j\} \rangle \neq V(G)$  for every  $a_j \in A$ , because if  $S_j = V(G)$ ,  $\bigcap_{a_i \in A'} \langle A' \setminus \{a_i\} \rangle = \emptyset$  for  $A' = A \setminus \{a_j\}$  then  $S_j$  is an edge of  $G$ . This implies that  $|A| = 3$  and  $G[A]$  is isomorphic to  $K_3$  and  $A$  is a convex set, a contradiction.

Then,  $h(G) = 2$ . ■

**Theorem 4.** *If a graph  $G$  contains a universal vertex, then  $h(G) = 2$ .*

*Proof.* Let  $G$  be a graph and  $v_1, \dots, v_k$ ,  $k \geq 1$ , universal vertices in  $G$ .

Let  $A$  be a subset of  $V(G)$ , where  $|A| = 3$ , and  $\mathcal{C}$  be the family of the convex sets containing at least two vertices of  $A$ . This implies that each  $S_i \in \mathcal{C}$  has at least two distinct vertices, so every convex set  $S_i$  contains

all the universal vertices of  $G$ , hence they belong to the intersection of the convex sets  $S_i$ . Then, by Theorem 1,  $G$  is 2-Helly. Since  $G$  is not trivial graph, the result follows.

Hence,  $h(G) = 2$ . ■

A graph is a *threshold graph* if there is a real number  $r$  and for every vertex  $v$  there is a real weight  $a_v$  such that:  $vw$  is an edge if and only if  $a_v + a_w \geq r$ .

**Corollary 5.** *If  $G$  is a threshold graph, then  $h(G) = 2$ .*

This result is a direct consequence of Theorem 4.

## 2.1 Path Graphs

The Helly number in a path  $P_n$  is directly related to the order of the graph.

**Theorem 6.** *If  $G = P_n$  is a path, then  $h(P_n) = \lceil \frac{2}{3}n \rceil$ .*

*Proof.* Let  $G = P_n$  be a graph with  $n$  vertices. We will label the vertices of  $G$  as follows:  $w_1, w_2, \dots, w_n$ , where  $d(w_1) = d(w_n) = 1$  and  $d(w_i) = 2$ , for  $i = 2, \dots, n - 1$ ; thus,  $w_i$  is adjacent to the vertices  $w_{i-1}$  and  $w_{i+1}$ .

Taking  $V_k = \{v_1, v_2, v_3, \dots, v_k\}$ , where  $v_j = w_i$ , for  $j = \lfloor \frac{2}{3}i \rfloor + 1$ , and  $i \not\equiv 0 \pmod{3}$ . Thus, given any three vertices in  $V_k$ , such vertices do not induce a subgraph  $P_3$ , and since in every  $P_3$  induced in  $G$  there are exactly two vertices of  $V_k$ , then  $V_k$  has maximum cardinality.

By Lemma 2 there exists in  $G$  a family  $\mathcal{C}_G$  of convex sets,  $(k - 1)$ -intersecting, with the empty core, i.e., by Theorem 1, the family  $\mathcal{C}_G = \{S_1, \dots, S_k\}$  is such that:

$$S_1 \supseteq \{v_2, v_3, \dots, v_k\} \not\supseteq \{v_1\}, S_2 \supseteq \{v_1, v_3, \dots, v_k\} \not\supseteq \{v_2\}, \dots, S_k \supseteq \{v_1, v_2, v_3, \dots, v_k\} \not\supseteq \{v_k\} \text{ and } \bigcap_{j=1}^k S_j = \emptyset.$$

It is easy to see that the inclusion of any other vertex of  $V(G)$  in  $V_k$  would imply, for some  $p$ , in the vertex sequence  $v_{p-1}$ ,  $v_p$  and  $v_{p+1}$ , that is, it would imply  $S_p \supseteq \{v_p\}$ , a contradiction. In general, if every induced

subgraph  $P_3$  in  $G$  has exactly two vertices of  $V_k$ , we have that  $|V_k|$  is maximum. Thus,  $h(G) = k$ .

In fact, for every three vertices in  $V(G)$ , exactly two vertices belong to  $V_k$ , then we considered three cases as a function of the number of vertices of  $G$ . Then we have the cases  $n \equiv 0 \pmod{3}$ ,  $n \equiv 1 \pmod{3}$  and  $n \equiv 2 \pmod{3}$ .

**Case 1:**  $n \equiv 0 \pmod{3}$ .

Clearly, for every three vertices  $w_p$ ,  $w_{p+1}$  and  $w_{p+2}$  in  $G$ , with  $p \equiv 1 \pmod{3}$ , we have exactly the first two of these vertices in  $V_k$  ( $w_p$  and  $w_{p+1}$ ), and in particular,  $v_k = w_{n-1}$ , i.e.,  $k = \frac{2}{3}n$ .

**Case 2:**  $n \equiv 1 \pmod{3}$ .

In this case,  $w_1$  and  $w_n$ , vertices of degree one, belong to  $V_k$  and  $w_{n-1} \notin V_k$ . Thus,  $k = \frac{2}{3}(n-1) + 1 = \frac{2}{3}n + \frac{1}{3}$ ,  $n \in \mathbb{N}$ , then  $k = \lceil \frac{2}{3}n \rceil$ .

**Case 3:**  $n \equiv 2 \pmod{3}$ .

In this case,  $w_{n-1}$  and  $w_n$  belong to  $V_k$ . Then,  $k = \frac{2}{3}(n-2) + 2 = \frac{2}{3}n + \frac{2}{3}$ , since  $k, n \in \mathbb{N}$ , then  $k = \lceil \frac{2}{3}n \rceil$ .

Since for  $n \equiv 0 \pmod{3}$  we have  $\frac{2}{3}n = \lceil \frac{2}{3}n \rceil$ , then  $k = \lceil \frac{2}{3}n \rceil$ , for  $n \in \mathbb{N}$ .

Hence,  $h(P_n) = \lceil \frac{2}{3}n \rceil$ . ■

## 2.2 Cycles

**Theorem 7.** *If  $G = C_n$  is a cycle, then  $h(C_n) = \lfloor \frac{2}{3}n \rfloor$ .*

*Proof.* Let  $G$  be a cycle  $C_n$  with  $n$  vertices. We will label the  $n$  vertices as  $w_1, w_2, \dots, w_n$ , where  $w_1$  and  $w_n$  are adjacent vertices and for each vertex  $w_i$ , for  $i = 2, \dots, n-1$ ;  $w_i$  is adjacent to the vertices  $w_{i-1}$  and  $w_{i+1}$ .

For the proof we consider the cases:  $n \equiv 0 \pmod{3}$ ,  $n \equiv 1 \pmod{3}$  and  $n \equiv 2 \pmod{3}$ .

**Case 1:**  $n \equiv 0 \pmod{3}$

Take  $V_k = \{v_1, v_2, \dots, v_k\}$ , where  $v_j = w_i$ , for  $j = \lfloor \frac{2}{3}i \rfloor + 1$ , and  $i \not\equiv 0 \pmod{3}$ . Thus,  $V_k$  has maximum cardinality, since every induced subgraph  $P_3$  in  $G$  has exactly two vertices of  $V_k$ . Then, by Lemma 2, there

exists a  $(k-1)$ -intersecting family  $\mathcal{C}_G$  containing  $k$  convex sets with empty core, which implies, by Theorem 1, that the family  $\mathcal{C}_G = \{S_1, \dots, S_k\}$  is such that:

$$S_1 \supseteq \{v_2, v_3, \dots, v_k\} \not\supseteq \{v_1\}, S_2 \supseteq \{v_1, v_3, \dots, v_k\} \not\supseteq \{v_2\}, \dots, S_k \supseteq \{v_1, v_2, v_3, \dots, v_k\} \not\supseteq \{v_k\} \text{ and } \bigcap_{j=1}^k S_j = \emptyset.$$

Hence,  $h(G) \geq k$ . Since  $|V_k|$  is maximum, then  $h(G) = k$ .

Thus, for every three vertices  $w_p, w_{p+1}$  and  $w_{p+2}$  in  $V(G)$ , with  $p \equiv 1 \pmod{3}$ , we will have exactly the first and two of these vertices in  $V_k$  ( $w_p$  and  $w_{p+1}$ ), and in particular,  $v_k = w_{n-1}$  and  $w_n \notin V_k$ , i.e.,  $k = \frac{2}{3}n$ .

**Case 2:**  $n \equiv 1 \pmod{3}$

Since  $n \equiv 1 \pmod{3}$ ,  $w_{n-1} \notin V_k$  and  $w_n \in V_k$ , then  $G[\{w_n, w_1, w_2\}] = P_3$ , a contradiction. Therefore,  $w_n$  cannot belong to  $V_k$ . Take  $V_k = \{v_1, v_2, \dots, v_k\}$ , where  $v_j = w_i$ , for  $j = \lfloor \frac{2}{3}i \rfloor + 1$ ,  $i \not\equiv 0 \pmod{3}$  and  $i < n$ . In this way,  $V_k$  has maximum cardinality, since every induced subgraph  $P_3$  in  $G \setminus \{w_n\}$  has exactly two vertices of  $V_k$  and for every vertex  $w_t \in V(G) \setminus V_k$ , there will exist in the set  $V_k \cup \{w_t\}$  a  $P_3$  induced subgraph, then  $|V_k|$  has maximum cardinality.

By Lemma 2 there exists in  $G$  a family  $\mathcal{C}_G$  of  $k$  convex sets,  $(k-1)$ -intersecting, with the empty core, then  $h(G) \geq k$ . Since  $|V_k|$  is maximum, then  $h(G) = k$ .

Thus,  $k = \frac{2}{3}(n-1) = \frac{2}{3}n - \frac{2}{3}$ , and since  $k, n \in \mathbb{N}$ , we have that  $k = \lfloor \frac{2}{3}n \rfloor$ .

**Case 3:**  $n \equiv 2 \pmod{3}$

Since  $w_1, w_2 \in V_k$  and  $w_{n-1} \in V_k$ , if  $w_n \in V_k$ , then  $G[\{w_{n-1}, w_n, w_1, w_2\}] = P_4$ , a contradiction. Hence,  $w_n$  cannot belong to  $V_k$ . Take  $V_k = \{v_1, v_2, \dots, v_k\}$ , where  $v_j = w_i$ , for  $j = \lfloor \frac{2}{3}i \rfloor + 1$ ,  $i \not\equiv 0 \pmod{3}$  and  $i < n$ . In this way,  $V_k$  has maximum cardinality, since every induced subgraph  $P_3$  in  $G \setminus \{w_n\}$  has exactly two vertices of  $V_k$  and for every vertex  $w_t \in V(G) \setminus V_k$ , there will exist in the set  $V_k \cup \{w_t\}$  a subgraph  $P_3$  induced, then  $|V_k|$  has maximum cardinality.

By Lemma 2, there exists in  $G$  a family  $\mathcal{C}_G$  of  $k$  convex sets,  $(k-1)$ -intersecting, with the empty core, then  $h(G) \geq k$ . Since  $|V_k|$  is maximum,

we have that  $h(G) = k$ .

Thus,  $k = \frac{2}{3}(n - 2) + 1 = \frac{2}{3}n - \frac{1}{3}$ , such as  $k, n \in \mathbb{N}$ , we have that  $k = \lfloor \frac{2}{3}n \rfloor$ .

Since for  $n \equiv 0 \pmod{3}$ , we have that  $\frac{2}{3}n = \lfloor \frac{2}{3}n \rfloor$ , thus  $k = \lfloor \frac{2}{3}n \rfloor$ , for  $n \in \mathbb{N}$ .

Hence,  $h(C_n) = \lfloor \frac{2}{3}n \rfloor$ . ■

### 2.3 Complete $k$ -partite graphs

A graph  $G$  is  $k$ -partite if  $V(G)$  can be partitioned in  $k$  independent sets and every possible edge that could connect vertices in different subsets is part of the graph.

**Theorem 8.** *If  $G$  is a complete  $k$ -partite graph, then  $h(G) = 2$ .*

*Proof.* Let  $G$  be a complete  $k$ -partite graph and  $S$  a proper subset of  $V(G)$  with at least 2 vertices. It is clear that if  $k = 2$ , then  $S$  is a convex set if and only if  $S$  is an edge, and if  $k \neq 2$ , then  $S$  is not a convex set, hence, by Lemma 3, the result follows. ■

**Corollary 9.** *If  $G = K_n$  is a complete graph, then  $h(K_n) = 2$ .*

Since complete graphs  $K_n$  are also complete  $k$ -partite graphs, such that each independent set contains only one vertex, then  $h(K_n) = 2$ .

## 3 Bounds

The bounds in Theorems 10 and 11 hold for all graph convexities [2].

**Theorem 10** ([2]). *Let  $G$  be a graph. Then  $h(G) \leq |V(G)|$ .*

If  $S$  is a convex set in a graph  $G$ ,  $h(G[S])$  is a lower bound of  $h(G)$ .

**Theorem 11** ([2]). *Let  $S$  be a convex set of a graph  $G$ . Then  $h(G[S]) \leq h(G)$ .*

**Definition 12.** A  $(k, l)$ -stable set  $\mathcal{S}$  is a set of vertices in a graph where:

- a) The elements of  $\mathcal{S}$  induces maximal cliques of the size less or equal to  $k$ .
- b) If  $u$  and  $v$  are two vertices in distinct cliques of  $\mathcal{S}$ ,  $d(u, v) \geq l$ .

**Corollary 13.** If  $\mathcal{S}$  is a maximum  $(2, 3)$ -stable set of  $G$ , then,  $h(G) \geq |\mathcal{S}|$ .

*Proof.* Let  $\mathcal{S}$  be a maximum  $(2, 3)$ -stable set with  $k$  vertices, such that  $\mathcal{S} = \{v_1, v_2, \dots, v_k\}$ .

The family of the convex sets  $\mathcal{C}_G = \{S_1, S_2, \dots, S_k\}$ :

$S_1 = \{v_2, v_3, \dots, v_k\}$ ,  $S_2 = \{v_1, v_3, \dots, v_k\}$ ,  $\dots$ ,  $S_{k-1} = \{v_1, \dots, v_{k-2}, v_k\}$ ,  
 $S_k = \{v_1, v_2, \dots, v_{k-1}\}$  is  $k$ -intersecting with empty core, i.e.,  $\bigcap_{i=1}^k S_i = \emptyset$ .

Thus,  $G$  is not  $(k - 1)$ -Helly. Since  $|\mathcal{S}|$  is maximum, the result follows. Hence,  $h(G) \geq |\mathcal{S}|$ . ■

## References

- [1] C. Berge, P. Duchet, A generalization of Gilmore's Theorem, *Recent Advances in Graph Theory* (Fielder, M., ed.), Acad. Praha, Prague, 1975, 49–55.
- [2] M. T. Carvalho Jr., O Número de Helly na Convexidade Geodética em Grafos, *Ph.D. Thesis*, UFRJ/COPPE, 2016.
- [3] C. C. Centeno, M. C. Dourado, Szwarcfiter J. L., On the convexity of paths of length two in undirected graphs, *Electronic Notes in Discrete Mathematics*, v. 32, pp. 15–18, 2009.
- [4] P. Duchet, Convex set in graphs II. Minimal path convexity. *Journal of Combinatorial Theory*, séries **B** 44, 1988, 307–316.
- [5] E. Helly, Ueber Mengen konvexer Koerper mit gemeinschaftlichen Punkter, Jahresber, *Math Verein.* v.32, 1923, 175–176.



- [6] M. L. J. Van de Vel, Theory of Convex Structures, *North-Holland Mathematical Library*, Elsevier, Amsterdam, London, New York, Tokyo, 1993.

COPPE - UFRJ,  
Rio de Janeiro - RJ - Brasil  
jayme@cos.ufrj.br  
IM - UFRJ,  
Rio de Janeiro - RJ - Brasil  
mitre@dcc.ufrj.br  
IME - UFF,  
Niterói - RJ - Brasil  
moisesteles@cos.ufrj.br