

A family of vector bundles on \mathbb{P}^3 of homological dimension 2 and $\chi(\text{End}E) = 1$

Sérgio Mendes, Rosa M. Miró-Roig, Helena Soares

Abstract

For any odd integer $\gamma \leq -5$, we construct a family of rank 3 vector bundles $\{E_\gamma\}$ on \mathbb{P}^3 with minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-2)^{\frac{3\gamma^2+8\gamma-3}{8}} \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{\frac{3\gamma^2+2\gamma-17}{4}} \rightarrow \mathcal{O}_{\mathbb{P}^n}^{\frac{3\gamma^2-4\gamma-7}{8}} \rightarrow E_\gamma \rightarrow 0.$$

and satisfying $\chi(\text{End}E_\gamma) = 1$.

1 Introduction

Notation 1.1. If E is a vector bundle on \mathbb{P}^n we denote

$$h^q(E) = \dim H^q(E) = \dim H^q(\mathbb{P}^n, E).$$

The Euler characteristic of E is defined by the integer

$$\chi(E) = \sum_{i=0}^n (-1)^i h^i(E).$$

Recall furthermore that we have an isomorphism $\text{End}E \cong E^\vee \otimes E$.

2000 AMS Subject Classification: 22D25, 22D10, 46L80.

Key Words and Phrases: General linear group, tempered dual, base change, K-theory.

Sérgio Mendes was partially supported by FCT through CMA-UBI (project UID/MAT/00212/2019). Rosa M. Miró-Roig was partially supported by MTM2016-78623-P. Helena Soares was partially supported by CIMA – Centro de Investigação em Matemática e Aplicações, Universidade de Évora, project UID/MAT/04674/2019 (Fundação para a Ciência e Tecnologia)

If E is a vector bundle on \mathbb{P}^n , with $n \geq 2$, of homological dimension 1 and defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^a \rightarrow \mathcal{O}_{\mathbb{P}^n}^b \rightarrow E \rightarrow 0 \quad (1)$$

then $\chi(\text{End } E) = a^2 + b^2 - (n+1)ab$ is the Euler characteristic of $\text{End } E$. It is well-known (see e.g. [6], Lemma 2.2.3) that $\chi(\text{End } E) = 1$ if and only if $(a, b) = (u_k, u_{k+1})$, with $k \geq 1$, where $\{u_k\}_{k \geq 0}$ is the sequence defined recursively by

$$\begin{cases} u_0 = 0, \\ u_1 = 1, \\ u_{k+1} = (n+1)u_k - u_{k-1}. \end{cases} \quad (2)$$

Moreover, for each pair of the form $(a, b) = (u_k, u_{k+1})$, where $\{u_k\}_{k \geq 0}$ is as above, the existence of a vector bundle E defined by a resolution of type (1) and thus satisfying $\chi(\text{End } E) = 1$ is guaranteed, for all $n \geq 2$ ([6], Theorem 2.2.7). It is worth recalling that one of the reasons for studying vector bundles satisfying $\chi(\text{End } E) = 1$ is that they provide good candidates for exceptional bundles, important objects when studying stability.

If we look at the case of homological dimension 2 or, more precisely, to vector bundles E defined by a minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^n}^c \rightarrow E \rightarrow 0,$$

with $n \geq 3$, then the Euler characteristic of $\text{End } E$ is now given by the ternary quadratic form

$$\chi(\text{End } E) = a^2 + b^2 + c^2 - (n+1)ab - (n+1)bc + \binom{n+2}{2}ac.$$

Finding a general formula of all integer and positive solutions of the equation $\chi(\text{End } E) = 1$ is much harder, so we will restrict to the case when $n = 3$ and for which this equation becomes

$$a^2 + b^2 + c^2 - 4ab - 4bc + 10ac = 1. \quad (3)$$

Besides, we would also like to know which solutions (a, b, c) of (3) give rise to a vector bundle E of homological dimension 2 and linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0. \quad (4)$$

In [5], Proposition 4.3, the authors already proved the existence of a certain family of vector bundles defined this way and satisfying $\chi(\text{End}E) = 1$. We can restate the referred proposition as follows.

Proposition 1.2. *There exists a vector bundle E of homological dimension 2 and linear resolution*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0,$$

with $(a, b, c) = (u_s, u_{s+1}, 4u_{s+1} - 10u_s)$, for each $s \geq 1$, where $\{u_s\}_{s \geq 1}$ is the sequence (2).

In particular, $\chi(\text{End}E) = 1$.

In this note we construct a new family of vector bundles with the required properties corresponding to 3-uples (a, b, c) different from the one provided by the above proposition, that is coming from triples of the form $(a, b, c) = (u_s, u_{s+1}, 4u_{s+1} - 10u_s)$.

This construction is described in the next section and inspired the authors for a more general result in a work in progress in [4].

2 A family of rank 3 bundles on \mathbb{P}^3 of homological dimension 2 and $\chi(\text{End}E) = 1$

Let E be a vector bundle on \mathbb{P}^3 of homological dimension 2 and linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0, \quad (5)$$

satisfying $\chi(\text{End}E) = 1$, that is, such that a, b and c satisfy

$$a^2 + b^2 + c^2 - 4ab - 4bc + 10ac = 1.$$

So a, b and c are integer positive solutions of the Diophantine equation

$$X^2 + Y^2 + Z^2 - 4XY - 4YZ + 10XZ = 1. \tag{6}$$

We would like to find for which solutions (a, b, c) of (6) there is a vector bundle E defined by a linear resolution of the form (5) with Betti numbers a, b and c (so that E will satisfy $\chi(\text{End } E) = 1$).

According to [4], any solution (a, b, c) of (6) is a triple of integers of the form

$$\begin{aligned} a &= \frac{\alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 + 2\alpha\beta + 4\gamma\delta}{8}, \\ b &= \frac{\alpha^2 + 2\gamma^2 - 6\beta^2 - 12\delta^2}{4}, \\ c &= \frac{\alpha^2 + 2\gamma^2 - 2\beta^2 - 4\delta^2 - 2\alpha\beta - 4\gamma\delta}{8}, \end{aligned} \tag{7}$$

for some $\alpha, \beta, \gamma, \delta \in \mathbb{Z}$, with α even and $\alpha\delta - \beta\gamma = \pm 1$. In order to these integers to be the Betti numbers of E they furthermore must satisfy

$$\begin{cases} a > 0 & \Leftrightarrow (\alpha + \beta)^2 + 2(\gamma + \delta)^2 > 3(\beta^2 + 2\delta^2), \\ a \leq b & \Leftrightarrow (\alpha - \beta)^2 + 2(\gamma - \delta)^2 \geq 11(\beta^2 + 2\delta^2). \end{cases} \tag{8}$$

Note also that under this parameterisation we have

$$\text{rk } E = a - b + c = \beta^2 + 2\delta^2. \tag{9}$$

Recall (Proposition 4.3 in [5]) that the existence of such a family of vector bundles, $\{E_s\}_{s \geq 1}$, is proved in the case when (a, b, c) is a triple of the form $(u_s, u_{s+1}, 4u_{s+1} - 10u_s)$, for $s \geq 1$, where $\{u_s\}_{s \geq 1}$ is the sequence (2). When $s = 1$ we get $(a, b, c) = (1, 4, 6)$ and E is the rank 3 vector bundle on \mathbb{P}^3 with minimal linear resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2) \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^4 \rightarrow \mathcal{O}_{\mathbb{P}^3}^6 \rightarrow E \rightarrow 0.$$

In fact, this is the only rank 3 case among the set of triples of this form. Using (7)–(9) we can easily deduce that this triple $(1, 4, 6)$ corresponds to

4 solutions of type $(\alpha, \beta, \gamma, \delta)$: $(-4, 1, -3, 1)$, $(-4, 1, 3, -1)$, $(4, -1, -3, 1)$ and $(4, -1, 3, -1)$.

We will next see that the triple $(1, 4, 6)$ does not exhaust all rank 3 possible solutions. We will construct a family $\{E_\gamma\}_{\{\gamma \in \mathbb{Z} : \gamma \text{ odd}, \gamma \leq -5\}}$ of rank 3 vector bundles with minimal linear resolution of type (5) and such that $\chi(\text{End}E) = 1$.

Suppose first that a vector bundle E of rank 3 exists defined by a linear resolution of type (5). Since E is globally generated with $h^0(E) = c$ and $3 = \text{rk} E \leq c$, we can take 2 general global sections of E , $\sigma_1, \sigma_2 \in H^0(E)$, and consider the curve $C = (\sigma_1 \wedge \sigma_2)_0$ defined as the degeneracy locus in \mathbb{P}^3 where σ_1 and σ_2 are linearly dependent. We thus have a short exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}^2 \rightarrow E \rightarrow \mathcal{I}_C(c_1(E)) \rightarrow 0,$$

where $c_1(E) = b - 2a$ denotes the first Chern class of E . Taking cohomology of this sequence we get

$$0 \rightarrow H^0(\mathcal{O}_{\mathbb{P}^3}^2) \cong \mathbb{C}^2 \rightarrow H^0(E) \cong \mathbb{C}^c \rightarrow H^0(\mathcal{I}_C(c_1(E))) \rightarrow 0,$$

and so $H^0(\mathcal{I}_C(c_1(E))) \neq 0$. Hence C is contained in a surface of degree $c_1(E)$ and therefore, $c_1(E) = b - 2a > 0$. We have just proved the following.

Proposition 2.1. *If E is a rank 3 vector bundle on \mathbb{P}^3 of homological dimension 2 and minimal linear resolution*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^c \rightarrow E \rightarrow 0,$$

then it has positive first Chern class, i.e. $b - 2a > 0$.

Therefore, our restrictions (8) above can be improved:

$$\begin{cases} a > 0 & \Leftrightarrow (\alpha + \beta)^2 + 2(\gamma + \delta)^2 > 3(\beta^2 + 2\delta^2), \\ b - 2a > 0 & \Leftrightarrow -\alpha\beta - 2\gamma\delta > 2(\beta^2 + 2\delta^2). \end{cases} \quad (10)$$

Furthermore, from the exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}^2 \rightarrow E \rightarrow \mathcal{I}_C(b - 2a) \rightarrow 0,$$

where $C = (\sigma_1 \wedge \sigma_2)_0$, we get a diagram

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \downarrow & & \downarrow & \\
 & & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & \\
 & & & \downarrow & & \downarrow & \\
 & & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & \\
 & & & \downarrow & & \downarrow & \\
 0 \rightarrow & \mathcal{O}_{\mathbb{P}^3}^2 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^c & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-2} & \longrightarrow 0 \\
 & \parallel & & \downarrow & & \downarrow & \\
 0 \rightarrow & \mathcal{O}_{\mathbb{P}^3}^2 & \longrightarrow & E & \longrightarrow & \mathcal{I}_C(b - 2a) & \longrightarrow 0 \\
 & \downarrow & & \downarrow & & \downarrow & \\
 & 0 & & 0 & & 0 &
 \end{array}$$

So, in order to reach our goal we will first prove the existence of a curve C in \mathbb{P}^3 with a resolution of type

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^{c-2} \rightarrow \mathcal{I}_C(b - 2a) \rightarrow 0.$$

We will then show that this implies the existence of a vector bundle E defined by (5) and satisfying $\chi(\text{End } E) = 1$.

Given any positive integer p , denote

$$D_p = \frac{p(p+1)}{2}$$

and

$$d_p = \begin{cases} \frac{p(p+2)}{4}, & \text{if } p \text{ is even,} \\ \frac{(p+1)^2}{4}, & \text{if } p \text{ is odd.} \end{cases}$$

Proposition 2.2. *Given an odd integer $\gamma \leq -5$, let*

$$p = -\frac{3\gamma + 7}{2} \quad \text{and} \quad s = \frac{3\gamma^2 + 8\gamma - 3}{8}.$$

Then, there is a smooth connected curve $C_\gamma \subset \mathbb{P}^3$ of degree

$$\text{deg}(C_\gamma) = \binom{p+1}{2} - s,$$

arithmetic genus

$$p_a = \frac{(2p+3)(p-1)(p-2)}{6} - s(p-1),$$

of maximal rank and with a minimal resolution

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2-p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1-p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_{C_\gamma} \rightarrow 0.$$

Proof. Let $d = \binom{p+1}{2} - s$ and let us first prove that $d_p \leq d \leq D_p$. In fact, on the one hand

$$d = \binom{p+1}{2} - s = \frac{(p+1)p}{2} - s \leq \frac{(p+1)p}{2} = D_p.$$

On the other hand, if p is even, we have

$$\begin{aligned} d_p &= \frac{p(p+2)}{4} \leq \binom{p+1}{2} - s = d \\ \Leftrightarrow \frac{p(p+2)}{4} &\leq \frac{(p+1)p}{2} - s \\ \Leftrightarrow 4s &\leq p^2 \Leftrightarrow \gamma \leq -5 \end{aligned}$$

If p is odd, we similarly obtain

$$d_p = \frac{(p+1)^2}{4} \leq \binom{p+1}{2} - s \Leftrightarrow 4s \leq p^2 - 1 \Leftrightarrow \gamma \leq -7,$$

and observe that when $\gamma = -5$, we get $p = 4$ even. From our hypotheses on γ and p we may conclude that it also holds $d_p \leq d$.

Therefore, by Proposition 2.2 in [1], there exists a smooth connected curve C_γ of degree $d = \binom{p+1}{2} - s$ such that

$$H^0(\mathbb{P}^3, \mathcal{I}_{C_\gamma}(p-1)) = H^1(\mathbb{P}^3, \mathcal{I}_{C_\gamma}(p-1)) = H^2(\mathbb{P}^3, \mathcal{I}_{C_\gamma}(p-1)) = 0.$$

In addition, by Remark 2.6 and Corollary 2.4 in [1], this curve has maximal rank and arithmetic genus

$$p_a = (p-1)d + 1 - \binom{p+2}{3} = \frac{(2p+3)(p-1)(p-2)}{6} - s(p-1).$$

Now, we may apply Corollary 5.1 in [2] to conclude that this curve C_γ has a p -linear resolution, that is, the ideal sheaf \mathcal{I}_C has a p -linear resolution, and that this resolution is of the following type:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2-p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1-p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_{C_\gamma} \rightarrow 0.$$

□

From now on we will concentrate on those solutions (a, b, c) of the diophantine equation (6) such that $\beta = \delta = 1$ and $\alpha = \gamma + 1$, with α an even integer (or equivalently, γ an odd integer).

After applying the required substitutions in (7), we get

$$a = \frac{3\gamma^2 + 8\gamma - 3}{8}, \quad b = \frac{3\gamma^2 + 2\gamma - 17}{4}, \quad c = \frac{3\gamma^2 - 4\gamma - 7}{8}.$$

Moreover, the inequalities (10) become

$$\begin{cases} a > 0 & \Leftrightarrow \frac{3\gamma^2 + 8\gamma - 3}{8} > 0, \\ b - 2a > 0 & \Leftrightarrow \frac{-3\gamma - 7}{2} > 0 \end{cases}$$

and we get that γ is an odd integer less than or equal to -5 ($\gamma \leq -5$).

Observing that $b - 2a = \frac{-3\gamma - 7}{2}$ and $a = \frac{3\gamma^2 + 8\gamma - 3}{8}$ are the integers p and s of Proposition 2.2, we know that there is a curve C_γ whose ideal sheaf has a linear resolution of the form

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2-p)^s \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1-p)^{2s+p} \rightarrow \mathcal{O}_{\mathbb{P}^3}(-p)^{s+p+1} \rightarrow \mathcal{I}_{C_\gamma} \rightarrow 0,$$

or equivalently, of the form

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^3}(-2)^a \rightarrow \mathcal{O}_{\mathbb{P}^3}(-1)^b \rightarrow \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \rightarrow \mathcal{I}_{C_\gamma}(b-2a) \rightarrow 0.$$

Now, Proposition 5.2 in [3] guarantees that we can take a non-trivial element

$$\begin{aligned} e_\gamma \in \text{Ext}^1(\mathcal{I}_{C_\gamma}(b-2a), \mathcal{O}_{\mathbb{P}^3}) &\cong \text{H}^2(\mathbb{P}^3, \mathcal{I}_{C_\gamma}(b-2a-4)) \\ &\cong \text{H}^1(C, \mathcal{O}_{C_\gamma}(b-2a-4)) \cong \text{H}^0(C, \omega_{C_\gamma}(4+2a-b)), \end{aligned}$$

and so we have a diagram

$$\begin{array}{ccccccc} & & & & 0 & & \\ & & & & \downarrow & & \\ & & & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & & \\ & & & & \downarrow & & \\ & & & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & & \\ & & & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^c & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \longrightarrow 0 \\ & & & & \searrow \phi & & \downarrow \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & E_\gamma & \longrightarrow & \mathcal{I}_{C_\gamma}(b-2a) \longrightarrow 0. \\ & & \downarrow & & & & \downarrow \\ & & 0 & & & & 0 \end{array}$$

Applying the covariant functor $\text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, -)$ to the bottom row we get the short exact sequence

$$0 \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1}) \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, E_\gamma) \rightarrow \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{I}_{C_\gamma}(b-2a)) \rightarrow 0.$$

Taking the morphism $\phi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, \mathcal{I}_{C_\gamma}(b-2a))$ we may conclude that there is a non-trivial morphism $\varphi \in \text{Hom}(\mathcal{O}_{\mathbb{P}^3}^c, E_\gamma)$ that allows us to

complete the above diagram to the following commutative diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}_{\mathbb{P}^3}(-2)^a & = & \mathcal{O}_{\mathbb{P}^3}(-2)^a & & \\
 & & \downarrow & & \downarrow & & \\
 & & \mathcal{O}_{\mathbb{P}^3}(-1)^b & = & \mathcal{O}_{\mathbb{P}^3}(-1)^b & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^c & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{b-a+1} \longrightarrow 0 \\
 & & & & \downarrow \varphi & \dashrightarrow \phi & \downarrow \\
 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}^{c-b+a-1} & \longrightarrow & E_\gamma & \longrightarrow & \mathcal{I}_{C_\gamma}(b-2a) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

where the middle column is a resolution of E_γ . So, the following theorem is proved.

Theorem 2.3. *Let $\gamma \leq -5$ be any odd integer. Then, there is a rank 3 vector bundle E_γ of homological dimension 2 defined by a linear resolution of type*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-2)^{\frac{3\gamma^2+8\gamma-3}{8}} \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{\frac{3\gamma^2+2\gamma-17}{4}} \rightarrow \mathcal{O}_{\mathbb{P}^n}^{\frac{3\gamma^2-4\gamma-7}{8}} \rightarrow E_\gamma \rightarrow 0,$$

and such that $\chi(\text{End } E_\gamma) = 1$.

References

- [1] E. Ballico, G. Bolondi, Ph. Ellia, R. M. Miró-Roig, Curves of maximum genus in range A and stick-figures, *Trans. Amer. Math. Soc.* 349 (1997), 4589–4608.
- [2] D. Eisenbud and S. Goto, Linear free resolutions and minimal multiplicity, *J. Algebra* 88 (1984), 89–133.

- [3] R. Hartshorne, R. M. Miró-Roig, On the intersection of ACM curves in \mathbb{P}^3 , *J. Pure Appl. Algebra* 219 (2015), 3195–3213.
- [4] S. Mendes, R. M. Miró-Roig and H. Soares, Existence of bundles E on \mathbb{P}^3 of homological dimension 2 and $\chi(\text{End} E) = 1$, preprint (2019).
- [5] R. M. Miró-Roig, H. Soares, Exceptional bundles of homological dimension k , *Forum Math.* 29 (2017), 701–715.
- [6] H. Soares, Steiner vector bundles on algebraic varieties. Ph.D. thesis, University of Barcelona 2008.

Sérgio Mendes

ISCTE

Av. das Forças Armadas, 1649-026, Lisbon, Portugal

Email: sergio.mendes@iscte.pt

Rosa Maria Miró-Roig Gran Via de les Corts Catalanes, Barcelona, Spain

Email: miro@ub.edu

Helena Soares

ISCTE

Av. das Forças Armadas, 1649-026, Lisbon, Portugal

Email: helena.soares@iscte.pt