

EXACT SOLUTIONS OF HEAT AND MASS TRANSFER EQUATIONS

Andrei D. Polyanin  Alexei I. Zhurov* 

Abstract

We outline generalized separation of variables as applied to nonlinear second-order partial differential equations. In this context, we suggest an approach to constructing exact solutions of nonlinear PDEs. The approach involves searching for transformations that “reduce the dimensionality” of the equation. New families of exact solutions of 3D nonlinear elliptic and parabolic equations that govern processes of heat and mass transfer in inhomogeneous anisotropic media are described. Moreover, the approach makes it possible to construct exact solutions of nonlinear wave equations. We also present solutions for three families of equations with logarithmic heat sources; the solutions are obtained by nonlinear separation of variables.

Introduction

Heat and mass transfer phenomena in a medium (solid, liquid, or gas) at rest are governed by heat (diffusion) equations [1–4]. For a homogeneous and isotropic medium, the thermal diffusivity (diffusion coefficient) that occurs in these equations is constant in the entire domain under study [5–7] and the heat (diffusion) equation is a linear partial differential equation with constant coefficients. In anisotropic media, the thermal diffusivity (diffusion coefficient) depends on the heat (mass) transfer direction and, in inhomogeneous media, can depend on the coordinates and even on the temperature [8–11]. In the last case, the heat (diffusion) equation is nonlinear. Various authors suggested many different

*The authors were partially supported by a research grant from the Russian Foundation for Basic Research (Project No. 97-02-17648).

AMS Subject Classification: Primary 35K55, 35J60; Secondary 80A20, 35L70.

Key words and phrases: Exact solutions, nonlinear PDE, heat and mass transfer equations.

relations to approximate the dependence of the transfer coefficients on the temperature or concentration, including linear, power-law, and exponential (e.g., see [8, 10, 12, 13]).

Heat (mass) transfer can be complicated by sources or sinks, which are associated with various physicochemical mechanisms of absorption and release of heat (substance). In combustion theory and nonisothermal macrokinetics of complex chemical reactions [4, 14], it is not infrequent that the power of heat sources/sinks depends on the temperature, often nonlinearly, e.g., exponentially [14] or in accordance with a power law [15]. In mass transfer theory, the rate of volumetric chemical reaction is widely approximated by power-law dependences on the concentration; at the same time, exponential, logarithmic, and other dependences are also used.

Exact solutions of heat and mass transfer equations play an important role in forming a proper understanding of qualitative features of various thermal and diffusion processes. Exact solutions of nonlinear equations make it possible to look into the mechanism of intricate phenomena such as spatial localization of heat transfer, peaking regimes, multiplicity and absence of steady states under certain conditions, etc. Even those particular exact solutions of PDEs which do not have a clear physical interpretation can be used as test problems for checking the correctness and accuracy of various numerical, asymptotic, and approximate analytical methods. In addition, model equations and problems that allow exact solutions serve as a basis for developing new numerical, asymptotic, and approximate methods. These, in turn, permit one to study more complicated problems that have no analytical solution.

Three basic approaches are traditionally used to seek exact solutions of nonlinear differential equations: (i) search for traveling-wave solutions, (ii) search for self-similar solutions, and (iii) application of groups to search for symmetries of the equations. The method of nonlinear separation of variables outlined below includes the first two approaches as its special cases and, very often, allows finding exact solutions that cannot be obtained by application of groups. Except for special cases of partial differential equations, the precise connec-

tion between symmetries and separation of variables is not fully understood at present, what is especially true for nonlinear cases (e.g., see [15, page xx]).

1. Structure of Exact Solutions for Some Heat and Mass Transfer Equations

Prior to describing nonlinear separation of variables, we first briefly remind the procedure of searching for self-similar solutions and that of separation of variables for linear equations.

1.1. Self-similar Solutions

For simplicity we consider the one-dimensional case. Self-similar solutions of one-dimensional heat equations are solutions of the form [17, 18]

$$T(x, t) = t^\beta f\left(\frac{x}{t^\gamma}\right), \quad (1.1)$$

where β and γ are some constants. The unknown function $f(x/t^\gamma)$ is determined by an ordinary differential equation resulting from the substitution of solution (1.1) into the original PDE.

More generally, self-similar solutions are said to be solutions of the form

$$T(x, t) = \varphi(t) f\left(\frac{x}{\psi(t)}\right). \quad (1.2)$$

The functions $\varphi(t)$ and $\psi(t)$ are chosen for reasons of convenience in the specific problem.

For illustration, we consider a nonlinear problem describing unsteady heat transfer in a semiinfinite plate, $x \geq 0$, with thermal diffusivity depending on the temperature, $a = a(T)$. Initially, for $t \leq 0$, the plate has a uniform temperature T_i . For $t > 0$, a temperature T_s is maintained at the plate boundary $x = 0$. Thus, we have the following boundary value problem:

$$\frac{\partial T}{\partial t} = \frac{\partial}{\partial x} \left[a(T) \frac{\partial T}{\partial x} \right]; \quad T|_{t=0} = T_i, \quad T|_{x=0} = T_s, \quad T|_{x \rightarrow \infty} \rightarrow T_i. \quad (1.3)$$

This problem has been the subject of much investigation in heat conduction theory and seepage theory (e.g., see [4, 19]).

A solution of problem (1.3) is sought in the form $T = T(\omega)$, $\omega = x/\sqrt{t}$, thus resulting in the ODE

$$[a(T) T'_\omega]'_\omega + \frac{1}{2}\omega T'_\omega = 0; \quad T|_{\omega=0} = T_s, \quad T|_{\omega \rightarrow \infty} \rightarrow T_i. \quad (1.4)$$

Solutions of problem (1.4) have been obtained for linear dependence of a on T [2, 20, 21], hyperbolic approximation [3, 22], and power-law dependence [14, 23].

A detailed list of exact solutions to equations of the form

$$\frac{\partial T}{\partial t} = \frac{\partial}{\partial x} \left[a(T) \frac{\partial T}{\partial x} \right] + \Phi(T)$$

can be found in [19] for $\Phi = 0$ and [24] for $\Phi \neq 0$.

1.2. Separation of Variables in Linear Equations

Many linear PDEs can be solved by separation of variables. For illustration, we consider a linear second order PDE of the form

$$F\left(x, t, T, \frac{\partial T}{\partial x}, \frac{\partial T}{\partial t}, \frac{\partial^2 T}{\partial x^2}, \frac{\partial^2 T}{\partial x \partial t}, \frac{\partial^2 T}{\partial t^2}\right) = 0, \quad (1.5)$$

with two independent variables, x and t , and the unknown function $T = T(x, t)$.

The solution procedure involves several stages, which are outlined below.

1. At the first stage, one seeks a particular solution of the form

$$T(x, t) = \varphi(x) \psi(t). \quad (1.6)$$

After substituting solution (1.6) into Eq. (1.5), one rewrites, if possible, the equation so that its left-hand side depends only on x (involves x , φ , φ'_x , and φ''_{xx}) and the right-hand side depends only on t (involves t , ψ , ψ'_t , and ψ''_{tt}). The equality is possible only if both sides are equal to the same constant, k , called the separation variable. Thus, one obtains ODEs for $\varphi(x)$ and $\psi(t)$ which contain the parameter k .

This procedure is called separation of variables in linear equations.

2. At the second stage, the principle of linear superposition is used—a linear combination of exact solutions of a linear equation is also an exact solution of this equation.

The functions φ and ψ in solution (1.6) depend not only on x and t but also on the separation constant, $\varphi = \varphi(x; k)$ and $\psi = \psi(t; k)$. For various values k_1, k_2, \dots of k we obtain distinct particular solutions of Eq. (1.5),

$$T_1(x, t) = \varphi_1(x) \psi_1(t), \quad T_2(x, t) = \varphi_2(x) \psi_2(t), \quad \dots,$$

where $\varphi_i = \varphi(x; k_i)$ and $\psi_i = \psi(t; k_i)$, $i = 1, 2, \dots$. The spectrum of possible values of k can be established from the boundary conditions.

According to the principle of linear superposition, the sum

$$T(x, t) = A_1 \varphi_1(x) \psi_1(t) + A_2 \varphi_2(x) \psi_2(t) + \dots, \quad (1.7)$$

where A_1, A_2, \dots are arbitrary constants, is an exact solution of the original equation. Formally, all A_i 's can be set equal to 1, thus combining them with the φ_i 's.

3. The third stage serves to determine the spectrum of k from the boundary conditions when solving specific problems. Here we arrive at the Sturm–Liouville eigenvalue problem for φ . The arbitrary constants A_i can be determined from the initial conditions.

Remark. Note that a lot of linear equations of mathematical physics can also admit exact solutions of the form

$$T(x, t) = \vartheta(x) + \chi(t), \quad (1.8)$$

where $\vartheta(x)$ and $\chi(t)$ are determined by the corresponding ODEs after separating the variables.

Example. Consider the linear equation

$$\frac{\partial C}{\partial t} = \frac{\partial}{\partial x} \left[D(x) \frac{\partial C}{\partial x} \right] + U(x) \frac{\partial C}{\partial x} + K_0 C + \Phi(t)$$

that governs a convective mass transfer at a speed of $-U(x)$, provided that the diffusion coefficient $D(x)$ depends on the coordinate, a first order chemical reaction takes place, K_0C , and there is a volume absorption of substance with intensity depending on time, $\Phi(t)$. This equation admits solutions of the form (1.8) but does not have exact solutions of the form (1.6). However, the equation admits more complicated solutions of the form

$$C(x, t) = \vartheta(x) \chi_1(t) + \chi_2(t), \quad (1.9)$$

where $\chi_1(t) = \exp(K_0 t)$ and the function $\chi_2(t)$ is determined by the first order ODE $\chi_2' = K_0 \chi_2 + \Phi(t)$.

1.3. Separation of Variables in Nonlinear Equations

Just as linear PDEs, some nonlinear equations admit exact solutions of the form (1.6). In this case, the functions $\varphi(x)$ and $\psi(t)$ are determined by the ODEs obtained by substituting Eq. (1.6) into the original equation and followed by nonlinear separation of variables.

Example 1. The nonlinear heat equation

$$\frac{\partial T}{\partial t} = \frac{\partial}{\partial x} \left(\alpha T^n \frac{\partial T}{\partial x} \right) \quad (1.10)$$

with the thermal diffusivity αT^n , where α and n are constants, admits exact solutions of the form (1.6) [19].

There are also nonlinear PDEs that admit exact solutions of the form (1.8).

Example 2. The nonlinear heat equation

$$\frac{\partial T}{\partial t} = \frac{\partial}{\partial x} \left(\alpha e^{\beta T} \frac{\partial T}{\partial x} \right) \quad (1.11)$$

with the thermal diffusivity $\alpha e^{\beta T}$, where α and β are constants, admits exact solutions of the form (1.8) [19].

Below we consider generalized separation of variables in nonlinear equations. Some aspects of this approach were considered in [30].

1. Suppose that a nonlinear equation for $T(x, t)$ is obtained from a linear equation for $u(x, t)$ admitting exact solutions of the form (1.6) or (1.8) by a nonlinear change of variable

$$T = F(u), \quad (1.12)$$

where $F(u)$ is some function. Then the nonlinear equation admits exact solutions of the form

$$T(x, t) = F(u), \quad u = \varphi(x) \psi(t), \quad (1.13)$$

$$T(x, t) = F(u), \quad u = \vartheta(x) + \chi(t). \quad (1.14)$$

For example, the above self-similar solution to the equation of (1.3) can be represented in the form (1.13) with $\varphi(x) = x$ and $\psi(t) = t^{-1/2}$.

Most commonly, solutions of nonlinear equations are sought in the form of traveling waves,

$$T(x, t) = F(u), \quad u = x + \lambda t. \quad (1.15)$$

Such solutions are special cases of Eq. (1.14) with $\vartheta(x) = x$ and $\chi(t) = \lambda t$. Note that solution (1.15) can also be represented in the form (1.13),

$$T(x, t) = F_1(v), \quad v = e^{x+\lambda t} = e^x e^{\lambda t}, \quad F_1(v) = F(\ln v).$$

Similarly, solution (1.14) can be represented in the form (1.13) by setting $u = \ln v$ and denoting $F(u) = F_1(v)$.

Usually, the functions φ and ψ or ϑ and χ , as well as the “temperature profile” $F = F(u)$, occurring in Eqs. (1.13) and (1.14) can be determined in the following two ways:

- The profile $F = F(u)$ is determined by an ODE resulting from the original equation after appropriate φ and ψ (or ϑ and χ) have been chosen. The functions φ and ψ (or ϑ and χ) also are determined by ODEs. Self-similar solutions and some more complicated solutions can be found in this way.

- The profile $F = F(u)$ is prescribed a priori on the basis of some considerations (e.g., a solution of a simpler auxiliary equation can be used as the profile) so that the variables can be separated. This leads to ODEs for φ and ψ (or ϑ and χ).

Table 1 presents some specific nonlinear equations that allow exact solutions of the form (1.13) or (1.14). We do not consider here self-similar solutions with $\varphi(x) = x$ and travelling wave type solutions.

2. Suppose a nonlinear PDE for $T(x, t)$ is obtained from a linear PDE for $u(x, t)$ admitting exact solutions of the form (1.7) by a nonlinear change of variable $T = F(u)$. Then the nonlinear equation admits solutions of the form

$$T(x, t) = F(u), \quad u = \varphi_1(x) \psi_1(t) + \varphi_2(x) \psi_2(t) + \cdots. \quad (1.16)$$

The structural formula (1.16) may now be used as a basis for seeking exact solutions to nonlinear equations that cannot be reduced to linear PDEs. The profile $F(u)$ and the functions $\varphi_1(x), \varphi_2(x), \dots, \psi_1(t), \psi_2(t), \dots$ are to be determined. It should be noted that generally solutions of this form cannot be obtained by group methods.

It is worth mentioning that in [27] exact solutions of the form (1.16) with $F(u) = u$, $\psi_2 = 1$, and $\psi_i = 0$, $i \geq 3$, were sought for PDEs with quadratic nonlinearities. In [30] a quite general procedure for seeking exact solutions of equations with quadratic nonlinearities for $F(u) = u$ is presented. Solutions of the form (1.16) are a natural extension of equations considered in the cited papers.

In the analysis of specific equations, it is useful to try the following special cases of formula (1.16):

$$T(x, t) = F(u), \quad u = \varphi_1(x) \psi_1(t) + \psi_2(t), \quad (1.17)$$

$$T(x, t) = F(u), \quad u = \varphi_1(x) \psi_1(t) + \varphi_2(x). \quad (1.18)$$

Table 2 presents some nonlinear equations admitting solutions of the form (1.16). One can see that solutions of the form (1.17) are most frequent.

Table 1: Some nonlinear PDEs admitting (1.13) or (1.14) type solutions

Equation	Solution structure	References
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + b \left(\frac{\partial T}{\partial x} \right)^2$	$T = \varphi(x) + \psi(t);$ $T = \frac{a}{b} \ln u, \quad u = \varphi(x) + \psi(t)$	[25]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(T^m \frac{\partial T}{\partial x} \right)$	$T = \varphi(x) \psi(t)$	[16, 17]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(e^{\lambda T} \frac{\partial T}{\partial x} \right)$	$T = \varphi(x) + \psi(t)$	[16, 19, 26]
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + aT \ln T$	$T = \varphi(x) \psi(t)$	[16, 24]
$\frac{\partial T}{\partial t} = \frac{a}{x^n} \frac{\partial}{\partial x} \left(x^n \frac{\partial T}{\partial x} \right) + aT \ln T$	$T = \varphi(x) \psi(t)$	[16, 27]
$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = a e^T$	$T = -2 \ln u, \quad u = \varphi(x) + \psi(y)$	[14]
$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = a \sinh T$	$T = 2 \ln \frac{1+u}{1-u}, \quad u = \varphi(x) \psi(y)$	[28]
$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = aT \ln T$	$T = e^u, \quad u = \varphi(x) + \psi(y)$	[28]
$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = a \sin T$	$T = 4 \arctan u, \quad u = \varphi(x) \psi(y)$	[28]
$\frac{\partial}{\partial x} \left(a x^n \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b y^m \frac{\partial T}{\partial y} \right) = c T^k$	$T = f(u), \quad u = \varphi(x) + \psi(y)$	[29]
$\frac{\partial}{\partial x} \left(a e^{\lambda x} \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b e^{\beta y} \frac{\partial T}{\partial y} \right) = c e^{\gamma T}$	$T = f(u), \quad u = \varphi(x) + \psi(y)$	[29]
$\frac{\partial}{\partial x} \left(a x^n \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b e^{\beta y} \frac{\partial T}{\partial y} \right) = c e^{\gamma T}$	$T = f(u), \quad u = \varphi(x) + \psi(y)$	[29]
$\frac{\partial}{\partial x} \left(a T^n \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b T^m \frac{\partial T}{\partial y} \right) = 0$	$T = \varphi(x) \psi(y)$	[29]
$\frac{\partial}{\partial x} \left(a e^{\lambda T} \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b e^{\beta T} \frac{\partial T}{\partial y} \right) = 0$	$T = \varphi(x) + \psi(y)$	[25]
$\frac{\partial^2 T}{\partial t^2} = \frac{\partial^2 T}{\partial x^2} + a e^T$	$T = -2 \ln u, \quad u = \varphi(x) + \psi(t)$	[25]
$\frac{\partial^2 T}{\partial t^2} = \frac{\partial^2 T}{\partial x^2} + a \sinh T$	$T = 2 \ln \frac{1+u}{1-u}, \quad u = \varphi(x) \psi(t)$	[28]
$\frac{\partial^2 T}{\partial t^2} = \frac{\partial^2 T}{\partial x^2} + aT \ln T$	$T = e^u, \quad u = \varphi(x) + \psi(t)$	[28]
$\frac{\partial^2 T}{\partial t^2} = \frac{\partial^2 T}{\partial x^2} + a \sin T$	$T = 4 \arctan u, \quad u = \varphi(x) \psi(t)$	[28]
Here $a, b, c, k, m, n, \beta, \gamma$, and λ are constants.		

It is important to note that in principle the representation (1.16) permits one to find exact solutions of nonlinear equations derived from a separable linear equation by a nonlinear transformation $T = F(u)$.

3. Suppose now that a nonlinear equation for $T(x, t)$ is obtained from a linear equation for $u(x, t)$ by a more general nonlinear change of variable

Table 2: Some nonlinear PDEs admitting solutions of the form (1.16)

Equation	Solution structure	References
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + bT \frac{\partial T}{\partial x}$	$T = 1/u, u = \varphi(x)\theta(t) + \psi(x)$	[26]
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + b \left(\frac{\partial T}{\partial x} \right)^2 + c_1 T + c_0$	$T = \varphi(t)x^2 + \psi(t)x + \chi(t)$	[29]
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + b \left(\frac{\partial T}{\partial x} \right)^2 + c_2 T^2 + c_1 T$	$T = \varphi(t)\theta(x) + \psi(t),$ $\theta(x) = e^{\lambda x}, \theta(x) = \sin(\lambda x)$	[27]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(T^m \frac{\partial T}{\partial x} \right)$	$T = u^{1/m}, u = \varphi(t)x^2 + \psi(t)$	[4, 19]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(T^m \frac{\partial T}{\partial x} \right) + bT$	$T = u^{1/m}, u = \varphi(t)x^2 + \psi(t)$	[24–26, 30]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(T^m \frac{\partial T}{\partial x} \right) + bT^{m+1}$	$T = u^{1/m}, u = \varphi(t)\theta(x) + \psi(t)$	[16]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(T^m \frac{\partial T}{\partial x} \right) + bT^{1-m}$	$T = u^{1/m}, u = \varphi(t)x^2 + \psi(t)$	[31]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(e^T \frac{\partial T}{\partial x} \right) + be^T + c$	$T = \ln u, u = \varphi(t)\theta(x) + \psi(t),$ $\theta(x) = e^{\lambda x}, \theta(x) = \sin(\lambda x)$	[32]
$\frac{\partial T}{\partial t} = a \frac{\partial}{\partial x} \left(e^T \frac{\partial T}{\partial x} \right) + b + ce^{-T}$	$T = \ln u, u = \varphi(t)x^2 + \psi(t)x + \chi(t)$	[29]
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + aT \ln T + bT$	$T = e^u, u = \varphi(t)x + \psi(t);$ $T = e^u, u = \varphi(t)x^2 + \psi(t)$	[29]
$\frac{\partial T}{\partial t} = a \frac{\partial^2 T}{\partial x^2} + T(a \ln^2 T + b \ln T + c)$	$T = e^u, u = \varphi(t)\theta(x) + \psi(t),$ $\theta(x) = e^{\lambda x}, \theta(x) = \sin(\lambda x)$	[27]
$\frac{\partial T}{\partial t} = \frac{a}{x^n} \frac{\partial}{\partial x} \left(x^n \frac{\partial T}{\partial x} \right) + aT \ln T$	$T = e^u, u = \varphi(t)x^2 + \psi(t)$	[16]
Here $a, b, c, c_0, c_1, c_2, m, n$, and λ are constants.		

$T = g(x, t)F(u) + h(x, t)$. By narrowing the classes of the functions $g(x, t)$ and $h(x, t)$, one arrives at more simple dependences, which may be used as a basis for seeking exact solutions of nonlinear equations that cannot be reduced to linear equations.

We suggest below structural formulas that are generalizations of relations (1.17) and (1.18):

$$T(x, t) = g(t)F(u) + h(t), \quad u = \varphi_1(x)\psi_1(t) + \psi_2(t), \quad (1.19)$$

$$T(x, t) = g(x)F(u) + h(x), \quad u = \varphi_1(x)\psi_1(t) + \varphi_2(x). \quad (1.20)$$

In the special case $\varphi_1(x) = x$ and $\psi_2(t) = 0$, formula (1.19) corresponds to generalized self-similar solutions.

2. Exact Solutions of 3D Nonlinear Heat and Mass Transfer Equations

2.1. Nonlinear Separation of Variables

Consider the following class of m -dimensional PDEs:

$$\sum_{i=1}^m \frac{\partial}{\partial x_i} \left[p_i(x_i) \frac{\partial w}{\partial x_i} \right] = P[w], \quad (2.1)$$

where the $p_i(x_i)$ are some functions to be established below, x_1, \dots, x_m are independent variables ($m \geq 2$). In general, the right-hand side of Eq. (2.1) is assumed to be a given nonlinear differential operator that depends on w , its derivatives with respect to independent variables x_{m+1}, \dots, x_k that do not enter the left-hand side, and the variables x_{m+1}, \dots, x_k themselves. The unknown w can play the role of temperature, concentration, or some other quantity.

We look for particular solutions of Eq. (2.1) of the form

$$w = w(r; \dots), \quad r^2 = \sum_{i=1}^m \varphi_i(x_i), \quad (2.2)$$

in which the number of independent variables is reduced by $m-1$. The unknown functions $\varphi_i(x_i)$ and $p_i(x_i)$ will be determined in the course of the study.

Substituting solution (2.2) into Eq. (2.1), we arrive at the equation

$$\frac{1}{4r^3} \left(r \frac{\partial^2 w}{\partial r^2} - \frac{\partial w}{\partial r} \right) \sum_{i=1}^m p_i(\varphi_i')^2 + \frac{1}{2r} \frac{\partial w}{\partial r} \sum_{i=1}^m (p_i \varphi_i')' = P[w], \quad (2.3)$$

where the primes denote the derivatives with respect to x_i .

The function of Eq. (2.2) is a solution of the original equation (2.1) only if the sums in Eq. (2.3) are constants or functions of r alone.

Generally, this is possible if

$$p_i(\varphi_i')^2 = A\varphi_i + A_i, \quad (p_i \varphi_i')' = B\varphi_i + B_i, \quad (2.4)$$

where A , A_i , B , and B_i are some constants ($i = 1, \dots, m$). In this case,

$$\sum_{i=1}^m p_i(\varphi_i')^2 = Ar^2 + A_\Sigma, \quad \sum_{i=1}^m (p_i \varphi_i')' = Br^2 + B_\Sigma, \quad A_\Sigma = \sum_{i=1}^m A_i, \quad B_\Sigma = \sum_{i=1}^m B_i.$$

For each i we have a system of two ODEs (2.4) for $p_i(x_i)$ and $\varphi_i(x_i)$.

Express p_i from the first equation in (2.4) in terms of φ_i to obtain

$$p_i = \frac{A\varphi_i + A_i}{(\varphi'_i)^2}. \quad (2.5)$$

Substituting this expression into the second equation in (2.4) yields the following autonomous equation for φ_i :

$$(A\varphi_i + A_i)\varphi''_i + (B\varphi_i + \mu_i)(\varphi'_i)^2 = 0, \quad (2.6)$$

where $\mu_i = B_i - A$. This equation can be solved by the substitution $\varphi'_i = z_i(\varphi_i)$.

For $A \neq 0$ the general solution of equation (2.6) can be represented in the implicit form

$$\begin{aligned} x_i + C_2 &= C_1 \int \exp\left(\frac{B\varphi_i}{A}\right) |A\varphi_i + A_i|^{\frac{A\mu_i - BA_i}{A^2}} d\varphi_i, \\ \varphi'_i = z_i(\varphi_i) &= \frac{1}{C_1} \exp\left(-\frac{B\varphi_i}{A}\right) |A\varphi_i + A_i|^{\frac{BA_i - A\mu_i}{A^2}}, \end{aligned} \quad (2.7)$$

where C_1 and C_2 are arbitrary constants.

For $A = 0$ and $A_i \neq 0$ the general solution of equation (2.6) is given by

$$\begin{aligned} x_i + C_2 &= C_1 \int \exp\left(\frac{B\varphi_i^2 + 2B_i\varphi_i}{2A_i}\right) d\varphi_i, \\ \varphi'_i = z_i(\varphi_i) &= \frac{1}{C_1} \exp\left(-\frac{B\varphi_i^2 + 2B_i\varphi_i}{2A_i}\right). \end{aligned} \quad (2.8)$$

In some cases the functions $p_i(x_i)$ and $\varphi_i(x_i)$ can be represented in explicit form. For example, if $A_i = B = B_i = 0$, from (2.7) and (2.5) we obtain

$$x_i + C_2 = \frac{C_1}{A} \ln |A\varphi_i|, \quad \varphi'_i = \frac{A}{C_1} \varphi_i, \quad p_i = \frac{A\varphi_i}{(\varphi'_i)^2}.$$

Whence,

$$p_i(x_i) = a_i e^{\lambda_i x_i}, \quad \varphi_i(x_i) = b_i e^{-\lambda_i x_i},$$

where $a_i = \pm C_1^2 e^{-AC_2/C_1}$, $\lambda_i = -A/C_1$, and $b_i = \pm A^{-1} e^{AC_2/C_1}$.

Table 3 shows special cases where the $p_i(x_i)$ and $\varphi_i(x_i)$ can be represented in explicit form.

On the basis of the preceding, we can formulate results for specific equations. In this paper, we confine ourselves to 3D equations and present exact solutions obtained using the above approach.

Table 3: Some cases where $p_i(x_i)$ and $\varphi_i(x_i)$ can be written out explicitly

#	$p_i(x_i)$	$\varphi_i(x_i)$	Relations for the parameters
1	$a_i x_i + s_i ^{n_i}$	$b_i x_i + s_i ^{2-n_i} + c_i$	$A_i = -Ac_i, B = 0,$ $B_i = \frac{A}{2-n_i}, b_i = \frac{A}{a_i(2-n_i)^2}$
2	$a_i e^{\lambda_i x_i}$	$b_i e^{-\lambda_i x_i} + c_i$	$A_i = -Ac_i, B = B_i = 0, b_i = \frac{A}{a_i \lambda_i^2}$
3	$a_i x_i^2$	$b_i \ln x_i + c_i$	$A = 0, A_i = a_i b_i^2, B = 0, B_i = a_i b_i$
4	$(a \ln x_i + b_i) x_i^2$	$c \ln x_i + d_i$	$A = ac, A_i = (b_i c - ad_i)c,$ $B = a, B_i = ac + (b_i c - ad_i)$

2.2. Exact Solutions of Heat/Mass Transfer and Wave Equations

Consider 3D equations corresponding to rows 1 and 2 in Table 3 which describe heat (mass) transfer or propagation of nonlinear waves in an anisotropic medium. In cases 1–4 below, we assume the operator $P[T]$ to be a nonlinear function $\Phi(T)$.

1. The equation ($k, m, n \neq 2$)

$$\frac{\partial}{\partial x} \left(a|x|^k \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(b|y|^m \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(c|z|^n \frac{\partial T}{\partial z} \right) = \Phi(T) \quad (2.9)$$

has exact solutions of the form ($A = \text{constant}$)

$$T = T(r), \quad r^2 = A \left[\frac{|x|^{2-k}}{a(2-k)^2} + \frac{|y|^{2-m}}{b(2-m)^2} + \frac{|z|^{2-n}}{c(2-n)^2} \right]. \quad (2.10)$$

The function $T(r)$ is determined by the ODE

$$T''_{rr} + \frac{D}{r} T'_r = \frac{4}{A} \Phi(T), \quad D = 2 \left(\frac{1}{2-k} + \frac{1}{2-m} + \frac{1}{2-n} \right) - 1. \quad (2.11)$$

This equation can be solved explicitly for $D = 1$ and $\Phi(T) = C \exp(\alpha T)$, where C and α are constants. For $D = 0$ and arbitrary $\Phi(T)$, Eq. (2.11) can be integrated in quadrature. For other exact solutions, see [33].

Note that $|x|$, $|y|$, and $|z|$ in Eqs. (2.9) and (2.10) can be replaced by $x + s_1$, $y + s_2$, and $z + s_3$, respectively, where s_1 , s_2 , and s_3 are arbitrary constants.

For $k = m = n = 0$ and $a = b = c$, Eq. (2.9) becomes a classical equation of heat (mass) transfer in an isotropic medium with heat release (volume reaction). In this case, solution (2.10), (2.11) corresponds to a spherically symmetric case.

2. The steady-state heat equation ($\lambda\mu\nu \neq 0$)

$$\frac{\partial}{\partial x} \left(ae^{\lambda x} \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(be^{\mu y} \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(ce^{\nu z} \frac{\partial T}{\partial z} \right) = \Phi(T) \quad (2.12)$$

admits solutions of the form

$$T = T(r), \quad r^2 = A \left(\frac{e^{-\lambda x}}{a\lambda^2} + \frac{e^{-\mu y}}{b\mu^2} + \frac{e^{-\nu z}}{c\nu^2} \right),$$

where $T(r)$ is determined by the ODE

$$T''_{rr} - \frac{1}{r} T'_r = \frac{4}{A} \Phi(T).$$

3. The equation ($n, m \neq 2$ and $\nu \neq 0$)

$$\frac{\partial}{\partial x} \left(ax^n \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(by^m \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(ce^{\nu z} \frac{\partial T}{\partial z} \right) = \Phi(T) \quad (2.13)$$

admits solutions of the form

$$T = T(r), \quad r^2 = A \left[\frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{e^{-\nu z}}{c\nu^2} \right],$$

where $T(r)$ is determined by the equation

$$T''_{rr} + \frac{D}{r} T'_r = \frac{4}{A} \Phi(T), \quad D = 2 \left(\frac{1}{2-n} + \frac{1}{2-m} \right) - 1.$$

4. The equation ($n \neq 2$ and $\mu\nu \neq 0$)

$$\frac{\partial}{\partial x} \left(ax^n \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(be^{\mu y} \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(ce^{\nu z} \frac{\partial T}{\partial z} \right) = \Phi(T) \quad (2.14)$$

has solutions of the form

$$T = T(r), \quad r^2 = A \left[\frac{x^{2-n}}{a(2-n)^2} + \frac{e^{-\mu y}}{b\mu^2} + \frac{e^{-\nu z}}{c\nu^2} \right].$$

The function $T(r)$ is determined by the ODE

$$T''_{rr} + \frac{D}{r} T'_r = \frac{4}{A} \Phi(T), \quad D = \frac{n}{2-n}.$$

For example, this equation is integrable in quadrature for $n = 0$ and arbitrary $\Phi(T)$ and explicitly for $n = 1$ and $\Phi(T) = Ce^{\alpha T}$, where C and α are constants.

5. Assume that $P[T] = \frac{\partial T}{\partial t} - \Phi(T)$. Consider the unsteady heat equation ($k, m, n \neq 2$)

$$\frac{\partial T}{\partial t} = \frac{\partial}{\partial x} \left(ax^k \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(by^m \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(cz^n \frac{\partial T}{\partial z} \right) + \Phi(T). \quad (2.15)$$

Following the approach of Subsection 2.1, we find that this equation has solutions of the form

$$T = T(t, r), \quad r^2 = 4A \left[\frac{x^{2-k}}{a(2-k)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{z^{2-n}}{c(2-n)^2} \right].$$

The function $T(t, r)$ satisfies a simpler PDE with two independent variables, specifically,

$$\frac{\partial T}{\partial t} = A \left(\frac{\partial^2 T}{\partial r^2} + \frac{D}{r} \frac{\partial T}{\partial r} \right) + \Phi(T), \quad D = \frac{2}{2-k} + \frac{2}{2-m} + \frac{2}{2-n} - 1.$$

For exact solutions of this equation, see [25].

Remark 1. Solutions of unsteady equations corresponding to Eqs. (2.12)–(2.14) can be constructed in a similar manner.

6. Assume that $P[T] = \partial^2 T / \partial t^2 - \Phi(T)$. Consider the following 3D equation describing the propagation of nonlinear waves in an inhomogeneous anisotropic medium ($\lambda\mu\nu \neq 0$):

$$\frac{\partial^2 T}{\partial t^2} = \frac{\partial}{\partial x} \left(ae^{\lambda x} \frac{\partial T}{\partial x} \right) + \frac{\partial}{\partial y} \left(be^{\mu y} \frac{\partial T}{\partial y} \right) + \frac{\partial}{\partial z} \left(ce^{\nu z} \frac{\partial T}{\partial z} \right) + \Phi(T). \quad (2.16)$$

It admits solutions of the form

$$T = T(r), \quad r^2 = A \left[-\frac{1}{4}(t+C)^2 + \frac{e^{-\lambda x}}{a\lambda^2} + \frac{e^{-\mu y}}{b\mu^2} + \frac{e^{-\nu z}}{c\nu^2} \right],$$

where A and C are arbitrary constants and $T(r)$ is determined by the ODE

$$T''_{rr} + \frac{4}{A} \Phi(T) = 0,$$

which is integrable in quadrature for any $\Phi(T)$.

Remark 2. Solutions of wave analogues of the heat equations (2.9), (2.13), and (2.14) can be constructed using similar considerations.

Remark 3. Two-dimensional analogues of the 3D equations considered above can be treated in a similar manner.

3. Nonlinear Equations with a Logarithmic Source

Following the method of nonlinear separation of variables outlined in Subsection 1.3, we found solutions of a number of other nonlinear equations. We chose to present three families of equations.

3.1. A 2D Steady-State Heat Equation

Consider the two-dimensional equation ($\alpha = \text{const}$, $\beta = \text{const}$)

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = \alpha T \ln \beta T. \quad (3.1)$$

1. This equation can be treated as a 2D special case of Eq. (2.9) with $m = n = 0$ and $\Phi(T) = \alpha T \ln \beta T$. Thus, Eq. (3.1) has solutions of the form

$$T = T(r), \quad T''_{rr} + \frac{1}{r}T'_r = \frac{\alpha}{A}T \ln \beta T, \quad r^2 = A[(x + C_1)^2 + (y + C_2)^2],$$

where A , C_1 , and C_2 are arbitrary constants.

2. Exact solutions of Eq. (3.1) can also be sought in the form $T = \frac{1}{\beta}e^{U(x,y)}$. With this change of variable, Eq. (3.1) becomes

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} + \left(\frac{\partial U}{\partial x}\right)^2 + \left(\frac{\partial U}{\partial y}\right)^2 = \alpha U. \quad (3.2)$$

Equation (3.2) admits traveling-wave solutions:

$$U(x, y) = F(u), \quad u = A_1 x + A_2 y + A_3, \quad (3.3)$$

where A_1 , A_2 , and A_3 are arbitrary constants. Substituting solution (3.3) into Eq. (3.2) and integrating the resulting equation, we obtain the dependence $F(u)$ in the implicit form

$$u = C_1 \pm \int \left[C_2 e^{-2F} + \frac{\alpha}{A_1^2 + A_2^2} \left(F - \frac{1}{2} \right) \right]^{-1/2} dF,$$

where C_1 and C_2 are arbitrary constants.

3. In addition, Eq. (3.2) has solutions of the form

$$U(x, y) = \varphi(x) + \psi(y).$$

Substituting this expression into Eq. (3.2) yields

$$\varphi''_{xx} + \varphi'^2_x - \alpha\varphi = -\psi''_{yy} - \psi'^2_y + \alpha\psi.$$

It follows that the variables separate and both sides must be equal to the same constant, which here can be set equal to zero. Solving the resulting equations, we obtain

$$x = A_1 \pm \int \left(B_1 e^{-2\varphi} + \alpha\varphi - \frac{1}{2}\alpha \right)^{-1/2} d\varphi, \quad (3.4)$$

$$y = A_2 \pm \int \left(B_2 e^{-2\psi} + \alpha\psi - \frac{1}{2}\alpha \right)^{-1/2} d\psi, \quad (3.5)$$

where A_1 , B_1 , A_2 , and B_2 are arbitrary constants.

4. Equation (3.2) admits also more complicated solutions of the form

$$U(x, y) = \varphi(\xi) + \psi(\eta), \quad \xi = x \cos \lambda - y \sin \lambda, \quad \eta = x \sin \lambda + y \cos \lambda,$$

where λ is an arbitrary constant and $\varphi(\xi)$ and $\psi(\eta)$ are determined by relations (3.4) and (3.5).

3.2. A 1D Unsteady Heat Equation

Consider the one-dimensional equation

$$\frac{\partial T}{\partial t} = \frac{a}{x^k} \frac{\partial}{\partial x} \left(x^k \frac{\partial T}{\partial x} \right) + f(t) T \ln T, \quad (3.6)$$

where a and k are some constants and $f(t)$ is an arbitrary function. Note that the values $k = 0, 1$, and 2 correspond to the plane, cylindrical, and spherical cases. The variables separate with the transformation

$$T(x, t) = e^{U(x, t)}, \quad U(x, t) = \varphi(t) x^2 + \psi(t).$$

Analysis shows that $\varphi(t)$ and $\psi(t)$ are determined by the following system of first order ODEs:

$$\varphi'_t = f\varphi + 4a\varphi^2, \quad \psi'_t = f\psi + 2a(k+1)\varphi.$$

The first, Bernoulli equation is integrable in quadrature for any $f = f(t)$. Whenever $\varphi(t)$ is found, the second, linear equation can be easily solved.

3.3. A 2D Unsteady Heat Equation

Consider the following two-dimensional heat equation:

$$\frac{\partial T}{\partial t} = a \left(\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} \right) - \alpha T \ln T.$$

We carry out the change of variable $T = e^{U(x,y,t)}$.

1. Exact solutions for U can be sought in the form $U(x, y, t) = \varphi(x, y) + \psi(t)$. The time-dependent term is expressed as $\psi(t) = Ae^{\alpha t}$, where A is an arbitrary constant. The function $\varphi(x, y)$ satisfies the steady-state equation

$$a \left(\frac{\partial^2 \varphi}{\partial x^2} + \frac{\partial^2 \varphi}{\partial y^2} \right) + a \left[\left(\frac{\partial \varphi}{\partial x} \right)^2 + \left(\frac{\partial \varphi}{\partial y} \right)^2 \right] - \alpha \varphi = 0,$$

which was considered in Subsection 3.1.

2. The equation for U admits other exact solutions, for example, $U(x, y, t) = \varphi(x, t) + \psi(y, t)$. The two unknown functions are determined by two independent one-dimensional nonlinear equations of the parabolic type,

$$\begin{aligned} \frac{\partial \varphi}{\partial t} &= a \frac{\partial^2 \varphi}{\partial x^2} + a \left(\frac{\partial \varphi}{\partial x} \right)^2 - \alpha \varphi, \\ \frac{\partial \psi}{\partial t} &= a \frac{\partial^2 \psi}{\partial y^2} + a \left(\frac{\partial \psi}{\partial y} \right)^2 - \alpha \psi. \end{aligned}$$

3. The following more sophisticated solutions are also possible:

$$U(x, y, t) = \varphi(\xi, t) + \psi(\eta, t), \quad \xi = x + \beta t, \quad \eta = y + \gamma t.$$

Here, β and γ are arbitrary constants. The unknown functions $\varphi(\xi, t)$ and $\psi(\eta, t)$ are determined by two independent one-dimensional nonlinear equations of the

parabolic type,

$$\begin{aligned}\frac{\partial \varphi}{\partial t} &= a \frac{\partial^2 \varphi}{\partial \xi^2} + a \left(\frac{\partial \varphi}{\partial \xi} \right)^2 - \beta \frac{\partial \varphi}{\partial \xi} - \alpha \varphi, \\ \frac{\partial \psi}{\partial t} &= a \frac{\partial^2 \psi}{\partial \eta^2} + a \left(\frac{\partial \psi}{\partial \eta} \right)^2 - \gamma \frac{\partial \psi}{\partial \eta} - \alpha \psi.\end{aligned}$$

To the special case $\varphi(\xi, t) = \varphi(\xi)$, $\psi(\eta, t) = \psi(\eta)$ there correspond autonomous ordinary differential equations.

4. Conclusions

Let us summarize the basic conclusions and results of the paper:

- The paper outlines the method of generalized separation of variables for nonlinear partial differential equations.
- In the context of this method, an approach is suggested which allows constructing exact solutions for some families of nonlinear PDEs. The approach is based on searching for transformations that reduce the equation to one with fewer independent variables.
- With this approach, new families of exact solutions of 3D nonlinear elliptic and parabolic equations that govern processes of heat and mass transfer in inhomogeneous anisotropic media are described.
- Exact solutions of some 3D hyperbolic equations describing the propagation of nonlinear waves in inhomogeneous anisotropic media are constructed.
- For three families of equations with logarithmic heat sources, a number of exact solutions are obtained by nonlinear separation of variables.

References

- [1] Carslaw, H. S.; Jaeger, J. C., *Conduction of Heat in Solids*, Clarendon Press, Oxford, (1984).

- [2] Kutateladze, S. S. *Foundations of Heat Transfer Theory* [in Russian], Atomizdat, Moscow, (1979).
- [3] Lykov, A. V., *Theory of Heat Conduction* [in Russian], Vysshaya Shkola, Moscow, (1967).
- [4] Zeldovich, Ya. B.; Barenblatt, G. I.; Librovich, V. B.; Makhviladze, G. M., *Mathematical Theory of Combustion and Explosion* [in Russian], Nauka, Moscow, (1980).
- [5] Gupalo, Yu. P.; Polyanin, A. D.; Ryazantsev, Yu. S., *Mass and Heat Exchange Between Reacting Particles and Flow* [in Russian], Nauka, Moscow, (1985).
- [6] Levich, V. G., *Physicochemical Hydrodynamics*, Prentice-Hall, Englewood Cliffs, New Jersey, (1962).
- [7] Clift, R.; Grace, J. R.; Weber, M. E., *Bubbles, Drops and Particles*, Acad. Press, New York, (1978).
- [8] Bretshnaider, S., *Properties of Fluids (Calculational Methods for Engineers)* [in Russian], Khimiya, Leningrad, (1966).
- [9] Polyanin, A. D.; Dilmán, V. V., *Methods of Modeling Equations and Analogies in Chemical Engineering*, CRC Press-Begell House, Boca Raton, (1994).
- [10] Reid, R. C.; Prausnitz, J. M.; Sherwood, T. K., *The Properties of Gases and Liquids*, McGraw-Hill Book Comp., New York, (1977).
- [11] Sherwood, T. K.; Pigford, R. L.; Wilkel, C. R., *Mass Transfer*, McGraw-Hill, New York, (1975).
- [12] Polyanin, A. D.; Kutepov, A. M.; Vyazmin, A. V.; Kazenin, D. A., *Hydrodynamics, Mass and Heat Transfer in Chemical Engineering*, Gordon & Breach Publ., Singapore, (2000).

- [13] Perry, J. H., *Chemical Engineers Handbook*, McGraw-Hill, New York, (1950).
- [14] Frank-Kamenetskii, D. A., *Diffusion and Heat Transfer in Chemical Kinetics* [in Russian], Nauka, Moscow, (1987).
- [15] Olver, P. J., *Application of Lie Groups to Differential Equations*, Springer-Verlag, New York, (1986).
- [16] Samarskii, A. A.; Galaktionov, V. A.; Kurdyumov, S. P.; Mikhailov, A. P., *Peaking Regimes in Problems for Quasilinear Parabolic Equations* [in Russian], Nauka, Moscow, (1987).
- [17] Barenblatt, G. I., *Dimensional Analysis*, Gordon and Breach Publ., New York, (1989).
- [18] Sedov, L. I., *Similarity and Dimensional Methods in Mechanics*, CRC Press, Boca Raton, (1993).
- [19] Ovsyannikov, L. V., *Group Analysis of Differential Equations*, Academic Press, New York, (1982).
- [20] Loitsyanskiy, L. G., *Laminar Boundary Layer* [in Russian], Fizmatgiz, Moscow, (1959).
- [21] Schlichting, H., *Boundary Layer Theory*, McGraw-Hill, New York, (1981).
- [22] Robillard, L., *On a series solution for the laminar boundary layer along a moving wall*, Trans. ASME, J. Appl. Mech., 38(2), (1978), p. 550.
- [23] Samarskii, A. A.; Sobol, I. M., *Examples of numerical analysis of temperature waves*, Zh. Vychisl. Mat. i Mat. Fiz., 3(4), (1963), p. 702.
- [24] Dorodnitsyn, V. A., *On invariant solutions of a nonlinear heat equation with a source*, Zh. Vychisl. Mat. i Mat. Fiz., 22(6), (1982), p. 1393.

- [25] Zaitsev, V. F.; Polyanin, A. D., *Handbook of Partial Differential Equations: Exact Solutions* [in Russian], MP Obrazovaniya, Moscow, (1996).
- [26] Ibragimov, N. H. (Editor), *CRC Handbook of Lie Group to Differential Equations*, Vol. 1, CRC Press, Boca Raton, (1994).
- [27] Galaktionov, V. A.; Posashkov, S. A., *On new exact solutions of parabolic equations with quadratic nonlinearities*, Zh. Vychisl. Mat. i Mat. Fiz., 29(4), (1989), p. 497.
- [28] Bullough, R. K.; Caudrey, P. J. (Editors), *Solitons*, Springer-Verlag, Berlin, (1980).
- [29] Polyanin, A. D.; Vyazmin, A. V.; Zhurov, A. I.; Kazenin, D. A., *Handbook of Exact Solutions of Heat and Mass Transfer Equations* [in Russian], Faktorial, Moscow, (1998).
- [30] Galaktionov, V. A.; Posashkov, S. A.; Svirshchevskii, S. R., *Generalized separation of variables for differential equations with polynomial right-hand sides*, Dif. Uravneniya, 31(2), (1995), p. 253.
- [31] Kersner, R., *On some properties of weak solutions of quasilinear degenerate parabolic equations*, Acta Math. Acad. Sci. Hung., 32(3–4), (1978), p. 301.
- [32] Berman, V. S.; Vostokov, V. V.; Ryazantsev, Yu. S., *On multiplicity of steady-state regimes with a chemical reaction*, Izv. AN SSSR, Mekh. Zhidkosti i Gaza [Fluid Dynamics], 3, (1982), p. 171.
- [33] Polyanin, A. D.; Zaitsev, V. F., *Handbook of Exact Solutions for Ordinary Differential Equations*, CRC Press, Boca Raton, New York, (1985).

Institute for Problems in Mechanics

Russian Academy of Sciences

101 Vernadsky Avenue

117526 Moscow, Russia

e-mail: polyanin@ipmnet.ru, zhurov@ipmnet.ru