

## A THEOREM OF REILLY FOR THE LINEARIZED OPERATOR OF $r^{\text{th}}$ MEAN CURVATURE AND APPLICATIONS TO STABILITY

H. Alencar  M. do Carmo H. Rosenberg 

In this note, we describe an estimate for the first eigenvalue of the linearized operator of the  $r^{\text{th}}$  mean curvature of immersed closed hypersurfaces of  $\mathbf{R}^{m+1}$  and  $\mathbf{H}^{m+1}$ ; we apply this to conclude that the stable closed hypersurfaces of  $\mathbf{R}^{m+1}$ , of constant curvature  $H_{r+1}$  are the round spheres.

**Reilly's Theorem:** Let  $M^n$  be a closed submanifold of  $\mathbf{R}^{m+1}$  and let  $H_1, \lambda_1$  be the mean curvature and first eigenvalue of the Laplacian of  $M$  respectively. R. Reilly proved [R1]:

$$\frac{\lambda_1}{n} \leq \frac{1}{\text{Vol}(M)} \int_M H_1^2 \quad (1)$$

and equality occurs precisely when  $M$  is minimally immersed in a sphere of  $\mathbf{R}^{m+1}$ . Hence when  $n = m$ , equality means  $M$  is a sphere.

Reilly's theorem extends easily to immersions in the unit sphere  $S^{m+1}$  by applying (1) to the immersion  $M \rightarrow S^{m+1} \subset \mathbf{R}^{m+2}$ :

$$\frac{\lambda_1}{n} - 1 \leq \frac{1}{\text{Vol}(M)} \int_M H_1^2. \quad (2)$$

For immersions of  $M^m$  in  $\mathbf{H}^{m+1}$ , the situation is more subtle. E. Heintze obtained some results [H] and the best result was obtained by A. El Soufi and S. Ilias [S-I]:

$$\frac{\lambda_1}{n} + 1 \leq \frac{1}{\text{Vol}(M)} \int_M H_1^2, m \geq 2, \quad (3)$$

and equality occurs precisely when  $M$  is minimally immersed in a geodesic sphere of radius  $\text{arch} \sqrt{\frac{m}{\lambda_1}}$ .

We pursue the study of  $\lambda_1 = \lambda_1(L_r)$ , where  $L_r$  is the linearised operator of  $S_{r+1} = \binom{m}{r+1} \mathbf{H}_{r+1}$  arising from normal variations of an immersed hypersurface  $M^m$  in  $\mathbf{R}^{m+1}$ . Here  $S_r$  is the  $r^{\text{th}}$  symmetric function of the eigenvalues of the shape operator  $A$ . Details concerning  $L_r$  can be found in [R2], [A-C-C], and [Ro]. Briefly,  $L_r(f) = \text{div}(T_r \nabla f)$ , where  $T_r$  is the  $r^{\text{th}}$  Newton transformation arising from  $A$ :

$$T_0 = I, T_r = S_r I - A T_{r-1} (\text{so } L_0 = \Delta).$$

**Our Theorem:** In  $\mathbf{R}^{m+1}$  we are able to generalize Reilly's result in the best possible way:

$$\lambda_1^{L_r} \int_M \mathbf{H}_r \leq C(r) \int_M \mathbf{H}_{r+1}^2, \quad (4)$$

where  $M$  is an immersed closed hypersurface in  $\mathbf{R}^{m+1}$  with  $\mathbf{H}_{r+1} > 0$  and  $C(r) = (m-r) \binom{m}{r}$ . Equality holds precisely when  $M$  is a sphere.

We also prove that if  $M$  extends to an isometric immersion of  $\Omega^{m+1} \rightarrow \mathbf{R}^{m+1}$ ,  $\partial\Omega = M$ , then

$$\lambda_1^{L_r} \leq \frac{C(r)}{(m+1)^2} \cdot \frac{V(M)}{V(\Omega)^2} \int_M \mathbf{H}_r$$

and equality holds precisely when  $M$  is a sphere.

Using this we prove such an immersion is  $r$ -stable if and only if  $M$  is a sphere. Here stability means  $M$  is a critical point of the functional  $\int_M \mathbf{H}_r + b\bar{V}(M)$  and the second derivative of this functional at  $M$  is non negative. Here  $b$  is a suitable constant and  $\bar{V}(M)$  is the (algebraic) volume bounded by  $M$ . This generalizes the theorems of Barbosa do Carmo [B-C] (stability of a constant mean curvature immersion means  $M$  is a sphere) and the theorem of Alencar, do Carmo and Colares [A-C-C] (for scalar curvature).

In  $\mathbf{H}^{m+1}$  we obtain an extrinsic upper bound for  $\lambda_1^{L_r}$  but it is not the best possible. Consequently our result does not yield stability here. The stability problem has been solved for scalar curvature in  $S^{m+1}$  [A-C-C] but this is not known in  $\mathbf{H}^{m+1}$ .

The techniques we use are properties of the operator  $L_r$  (when it is elliptic, formulae for  $L_r$  of particular functions), integral geometry, and inequalities involving the mean curvatures. Details will appear elsewhere.

## References

- [A-C-C] Alencar, H., do Carmo, M. and Colares, A.G., *Stable hypersurfaces with constant scalar curvature*, To appear Math. Z., 1993,
- [B-C] Barbosa, L. and do Carmo, M., *Stability of hypersurfaces with constant mean curvature*, Math. Z. **185** (1984), 339-353.
- [H] Heintze, E., *Extrinsic Upper Bounds for  $\lambda_1$* , Math. Ann. **28** (1988), 389-402.
- [R1] Reilly, R., *On the first eigenvalues of the Laplacian for compact submanifolds, of Euclidean space*, Comment. Math. Helv. **52** (1977), 525-533.
- [R2] Reilly, R., *Variational properties of functions of the mean curvatures for hypersurfaces in space forms*, J. Diff. Geom. **8** (1973), 465-477.
- [Ro] Rosenberg, H., *Hypersurfaces of Constant Curvature in Space Forms*, To appear Bull. Sc. Math. 1993.
- [S-I] El Soufi, A. and Ilias, S., *Une inegalite du type "Reilly" pour les sous-varietes de l'espace hyperbolique*, Comm. Math. Helv., **67** (1992), 167-181.

H. Alencar  
Universidade Federal de Alagoas  
Departamento de Matemática  
57080 Maceió, AL, Brasil

M. do Carmo  
IMPA  
Estrada Dona Castorina, 110  
22460 Rio de Janeiro, RJ, Brasil

H. Rosenberg

Department de Mathematiques

Université Paris VII

2 Place Jussieu

72251 Paris, Cedex 05, France