

Constructing pairs of Laplacian equienergetic threshold graphs

R. R. Del-Vecchio  G. B. Pereira

C. T. M. Vinagre 

Abstract

The Laplacian energy of a graph G on n vertices and m edges is defined as the sum of absolute values of the differences between each Laplacian eigenvalue of G and the average degree $2m/n$. In this work we construct pairs of threshold graphs of same order, with same Laplacian energy and different sets of Laplacian eigenvalues.

1 Introduction

Let G be a simple and undirect graph on vertices v_1, \dots, v_n . The *Laplacian matrix* $L = L(G)$ of G is the $n \times n$ matrix for which the entry L_{ii} is the degree of vertex v_i , $1 \leq i \leq n$, and the entries L_{ij} are -1 , if vertex v_i and v_j are adjacent in G , and 0 otherwise. The matrix L is symmetric, positive-semidefinite and always has 0 as an eigenvalue. For these and other properties of the L matrix, see [Mer94]. As usual, we denote the L -eigenvalues as $\mu_1, \mu_2, \dots, \mu_n$, where $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n = 0$; these real numbers constitute the *Laplacian spectrum* of G . Two graphs are

2000 AMS Subject Classification: : 05C50 and 05C69.

Key Words and Phrases: Laplacian energy of graph, threshold graph.

The first author was partially supported by CNPq.

cospectral if they share the same Laplacian spectrum, otherwise they are *non-cospectral*.

If G has m edges, then $2m/n$ is its *average degree* and the *Laplacian energy* of G is defined as $LE(G) = \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|$. After introduced by Gutman and Zhou in [GZ06], this concept has been extensively investigated. In particular, the problem of constructing families of non-cospectral graphs with same order and equal Laplacian energy (called *Laplacian-equienergetic graphs* or simply *L-equienergetic graphs*) has already been studied in [Ste09] and [FHT14], but a general characterization remains an open problem.

Our investigation concerns the construction of *L-equienergetic threshold graphs*.

Threshold graphs were introduced by Chvátal and Hammer [CH77] and independently, by Henderson and Zalcstein [HZ77], in 1977. They constitute an important class of graphs due to their numerous applications in diverse areas. Threshold graphs can be characterized in many ways. In this paper, a *threshold graph* is obtained through an iterative process which starts with an isolated vertex, and where, at each step, either a new isolated vertex is added, or a vertex adjacent to all previous vertices (*dominating vertex*) is added. Following this construction, a threshold graph can be represented by a string of 0's and 1's, corresponding respectively to isolated vertices and *dominating* vertices. The threshold graph constructed in Figure 1 has 0-1-string $(0, 0, 0, 1, 0, 0, 1)$.




Figure 1: Constructing the threshold graph with string $(0, 0, 0, 1, 0, 0, 1)$.

The number of characters 1 in the string (that is, the number of domi-

nating vertices in the graph) is called the *trace* of the graph and denoted by $T = T(G)$. The dominating vertices with the first vertex inserted in the graph constitute a clique. Thus, the clique number of the graph is equal to $T + 1$. For example, the graph in Figure 1 has clique number 3 ($= T + 1$). The other 4 ($= n - T - 1$) vertices constitute an independent set.

The 0-1-string of a threshold graph also provides its degree sequence $d = [d_1, d_2, \dots, d_i, \dots, d_n]$, where $d_1 \geq d_2 \geq \dots \geq d_n$. For example, the graph in Figure 1 has degree sequence $[6, 4, 2, 2, 2, 1, 1]$.

The following known result furnishes the sequence of L -eigenvalues of a threshold graph from its degree sequence, showing that all of them are integers.

Theorem 1.1 ([Mer94]). Let G be a threshold graph on n vertices, with trace T and degree sequence $[d_1, d_2, \dots, d_n]$ arranged in non increasing order. Then for the Laplacian eigenvalues of G it holds that $\mu_i = d_i + 1$, if $1 \leq i \leq T$, $\mu_i = d_{i+1}$, if $T + 1 \leq i \leq n - 1$, and $\mu_n = 0$.

In [VDVJT13], the authors establish an explicit formula to compute the Laplacian energy of threshold graphs satisfying certain hypothesis.

Theorem 1.2 ([VDVJT13]). Let G be a threshold graph on n vertices, m edges and trace T with $3 \leq T \leq n - 1$. If the Laplacian eigenvalues of G satisfy $\mu_T \geq 2m/n \geq \mu_{T+1}$ then the Laplacian energy of G is given by

$$LE(G) = T^2 + T + \left(1 - \frac{2T}{n}\right) 2m.$$

2 Families of threshold graphs satisfying conditions of Theorem 1.2

Given integers $n \geq 3$ and $T \geq 1$, the graph on n vertices obtained by attaching $n - T - 1$ pendent vertices to the same vertex of the complete graph K_{T+1} is said to be the *pineapple on n vertices and trace T* and

denoted $P_{n,T}$. Its 0-1 string is $(0, \underbrace{1, 1, \dots, 1}_{T-1}, \underbrace{0, 0, \dots, 0}_{n-T-1}, 1)$. For fixed n and T , it is the threshold graph with least number of edges. The graph $P_{n,T}$ has degree sequence $[n-1, T$ (T times), $1((n-T-1)$ times)], $\frac{1}{2}(T^2+T) + (n-T-1)$ edges and its Laplacian spectrum is $n, T+1$ ($(T-1)$ times), 1 ($(n-T-1)$ times) and 0 .

In what follows, n and T are integers such that $4 \leq T \leq n-2$.

2.1 A known family

In [VDVJT13], the authors exhibit a family of threshold graphs satisfying the hypothesis of Theorem 1.2. Denote by $\mathcal{G}_{n,T}$ the family of threshold graphs G_t on n vertices and trace T , where $G_0 = P_{n,T}$ and for $t, 1 \leq t \leq (T-1)(n-T-1)$, G_t is obtained from G_0 by the addition of t edges to specific vertices of G_0 in such way that the 0-1-strings of the graphs obtained are:

$$\begin{aligned}
 s_0 &= (0, \underbrace{1, \dots, 1}_{T-1}, \underbrace{0, 0, \dots, 0}_{n-T-1}, 1); & s_1 &= (0, \underbrace{1, \dots, 1}_{T-2}, 0, 1, 0, 0, \dots, 0, 1); \\
 s_2 &= (0, \underbrace{1, \dots, 1}_{T-2}, 0, 0, 1, 0, 0, \dots, 0, 1); & \dots & s_{(n-T-1)} = (0, \underbrace{1, \dots, 1}_{T-2}, 0, 0, \dots, 0, 1, 1); \\
 & & s_{(n-T-1)+1} &= (0, \underbrace{1, \dots, 1}_{T-3}, 0, 1, 0, 0, \dots, 0, 1, 1); & \dots \\
 s_{2(n-T-1)} &= (0, \underbrace{1, \dots, 1}_{T-3}, 0, \dots, 0, 1, 1, 1); & \dots & s_{(T-2)(n-T-1)} = (0, 1, 0, \dots, 0, \underbrace{1, \dots, 1}_{T-2}, 1, 1); \\
 & & s_{(T-2)(n-T-1)+1} &= (0, 0, 1, 0, \dots, 0, \underbrace{1, \dots, 1}_{T-2}, 1, 1); & \dots \\
 & & \dots & s_{(T-1)(n-T-1)} = (0, \dots, 0, \underbrace{1, \dots, 1}_{T-1}, 1).
 \end{aligned}$$

In [VDVJT13], conditions on the number of inserted edges t are given in order to identify the graphs G_t of $\mathcal{G}_{n,T}$ satisfying the hypothesis of Theorem 1.2. The introduction of the parameters t^b and t^\sharp , both depending on n and T , provides bounds on t in order to compute the Laplacian energy of these graphs.




Figure 2: The graphs $G_0 = P_{13,5}$, G_3 and G_9 of family $\mathcal{G}_{13,5}$.

Theorem 2.1 ([VDVJT13]). For fixed n and T , let $t^\flat = \frac{1}{2}(n-T)(T-1) + 1$ and $t^\sharp = \frac{1}{n-2} [(1-n)T^2 + (n^2-1)T + (3-2n)n - 2]$. Thus $1 \leq t^\flat \leq (n-T-1)(T-2) \leq t^\sharp \leq (n-T-1)(T-1)$. Furthermore, for each integer t , $1 \leq t \leq (n-T-1)(T-1)$, if $t \leq t^\flat$ or $t \geq t^\sharp$ then $LE(G_t) = T^2 + T + \left(2 - \frac{4T}{n}\right) m_t$, where m_t denotes the number of edges of the graph G_t of $\mathcal{G}_{n,t}$.

As pointed out in [VDVJT13], at least half of the threshold graphs in $\mathcal{G}_{n,T}$ satisfy the conditions of Theorem 2.1.

2.2 A new family

In this work, we present the construction of another family of threshold graphs for which Theorem 1.2 still holds.

The graphs F_t of the family $\mathcal{F}_{n,T}$ have n vertices and trace T . They are also obtained from the pineapple $F_0 = P_{n,T}$ by the insertion of t edges ($1 \leq t \leq (T-1)(n-T-1)$) in such way their respective 0-1-strings $s_1, \dots, s_t \dots s_{(n-T-1)(T-1)}$ are:

$$\begin{aligned}
 s_1 &= (0, \underbrace{1, \dots, 1}_{T-2}, 0, 1, 0, \dots, 0, 1); & s_2 &= (0, \underbrace{1, \dots, 1}_{T-3}, 0, 1, 1, 0, \dots, 0, 1); \\
 s_3 &= (0, \underbrace{1, \dots, 1}_{T-4}, 0, 1, 1, 1, 0, \dots, 0, 1); & \dots & s_{(T-1)} = (0, 0, \underbrace{1, \dots, 1}_{T-1}, 0, \dots, 0, 1); \\
 s_{(T-1)+1} &= (0, 0, \underbrace{1, \dots, 1}_{T-2}, 0, 1, 0, \dots, 0, 1); & s_{(T-1)+2} &= (0, 0, \underbrace{1, \dots, 1}_{T-3}, 0, 1, 1, 0, \dots, 0, 1); \\
 \dots & s_{2(T-1)} = (0, 0, 0, \underbrace{1, \dots, 1}_{T-1}, 0, \dots, 0, 1); & s_{2(T-1)+1} &= (0, 0, 0, \underbrace{1, \dots, 1}_{T-2}, 0, 1, 0, \dots, 0, 1);
 \end{aligned}$$

$$\cdots s_{(n-T-1)(T-1)-1} = (0, \dots, 0, 1, 0, \underbrace{1, \dots, 1}_{T-1}) \cdots$$

$$\cdots \dots s_{(n-T-1)(T-1)} = (0, \dots, 0, 1, 1, 1, \dots, 1, 1)$$




Figure 3: Graphs in family $\mathcal{F}_{8,4}$: $F_0 = P_{8,4}$, F_1 , F_2 , F_3 , F_4 and F_9 .

Reasonings analogous to Proposition 2 of [VDVJT13] we can prove that:

Theorem 2.2. For fixed n and T , let $t_1 = \frac{T(T-1)}{n-2} + 1$ and $t_2 = \frac{1}{2}(n-T)(T-1) + 1 - \frac{n}{2}$. Thus $1 \leq t_1 \leq (n-T-1)(T-2) \leq t_2 \leq (n-T-1)(n-1)$. Furthermore, for all t , $1 \leq t \leq (n-T-1)(T-1)$, if $1 \leq t \leq t_1$ or $t \geq t_2$ then $LE(F_t) = T^2 + T + \left(2 - \frac{4T}{n}\right) m_t$, where m_t denotes the number of edges of the graph F_t .

Proof: Analogously to the proof of Theorem 2.1 (Proposition 2 of [VDVJT13]), the three statements below must be proved:

1. $\mu_T \geq \frac{2m_t}{n}; \forall 1 \leq t \leq (n-T-1)(T-1)$
2. for $1 \leq t \leq (T-2)$, $\mu_{T+1} \leq \frac{2m_t}{n}$ if and only if $t \leq t_1$;
3. for $(T-1) \leq t \leq (n-T-1)(T-1)$, $\mu_{T+1} \leq \frac{2m_t}{n}$ if and only if $t \geq t_2$.

■

Numerical experiments show that at least half of the threshold graphs in $\mathcal{F}_{n,T}$ satisfy the conditions of Theorem 2.2.

3 Constructing pairs of L -equienergetic threshold graphs

Threshold graphs are determined by their L -spectra, that is, if G and H are two L -cospectral threshold graphs then they are isomorphic [Mer94].

In [Ste09], the author constructs large sets of L -equienergetic threshold graphs all of them having equal trace. Pairs of L -equienergetic threshold graphs with different traces are exhibited in [VDVJT13]. In what follows, we present infinite pairs of L -equienergetic threshold graphs.

In the next result, we present pairs of L -equienergetic graphs where both graphs were constructed by the same way although with different traces.

Theorem 3.1. For fixed n and T , let $T' = n - T$. For t such that $1 \leq t \leq (n - T - 1)(T - 1)$, let $s = (n - T' - 1)(T' - 1) - (t - 1)$. If $t \leq t_1(T)$ of Theorem 2.2 then $LE(F_t) = LE(F_s)$, where F_t belongs to family $\mathcal{F}_{n,T}$ and F_s to family $\mathcal{F}_{n,T'}$.

Proof: Firstly, we note that F_t and F_s have different spectra since their traces are distinct. By algebraic manipulations it can be verified that if $t \leq t_1(T)$ then $s \geq t_2(T')$ and so, $LE(F_t)$ and $LE(F_s)$ can be obtained by Theorem 2.2. Let m_t and m_s denote the number of edges of F_t and of F_s , respectively. Then

$$\begin{aligned} LE(F_t) &= T^2 + T + \left(1 - \frac{2T}{n}\right) 2m_t = \\ &= T^2 + T + \left(1 - \frac{2T}{n}\right) 2 \left(\frac{T(T+1)}{2} + n - T - 1 + t\right) = \\ &= 2 \left[T^2 - 2T + n - \frac{T^2}{n} (T - 1) + (t - 1) \left(1 - \frac{2T}{n}\right) \right]. \end{aligned}$$

$$\text{Since } 2m_s = 2 \left(\frac{T'(T'+1)}{2} + n - T' - 1 + (T' - 1)(n - T' - 1) - (t - 1) \right) =$$

$$= n^2 - T^2 + T - n + 2 - 2t$$

$$\text{then } \left(1 - \frac{2T'}{n}\right) 2m_s = \left(\frac{2T'}{n} - 1\right) (n^2 - T^2 + T - n + 2 - 2t) =$$

$$= T^2 + 2Tn - 3T - n^2 + n - \frac{2T^2}{n} (T - 1) + (t - 1) \left(2 - \frac{4T}{n}\right). \text{ Thus it follows that}$$

$$LE(F_s) = T'^2 + T' + \left(1 - \frac{2T'}{n}\right) 2m_s = 2 \left[T^2 - 2T + n - \frac{T^2}{n} (T - 1) + (t - 1) \left(1 - \frac{2T}{n}\right) \right] = LE(F_t).$$

■

In the sequence, we construct pairs of L -equienergetic graphs where one belongs to family $\mathcal{G}_{n,T}$ and the other, to family $\mathcal{F}_{n,T}$.

Theorem 3.2. Consider the parameters t^b , t^\sharp of Theorems 2.1 and t_1 and t_2 of Theorem 2.2. For all integer t with $1 \leq t \leq (n - T - 1)(T - 1)$, if $t \leq \min\{t^b, t_1\}$ or $t \geq \max\{t^\sharp, t_2\}$ then the graphs G_t in family $\mathcal{G}_{n,T}$ and F_t in family $\mathcal{F}_{n,T}$ are L -equienergetic.

Proof: As the graphs G_t and F_t have the same trace and same order n , the assertion follows from Theorems 2.1 and 2.2, since $LE(G_t) = T^2 + T + \left(1 - \frac{2T}{n}\right) 2m_t$ and $LE(F_t) = T^2 + T + \left(1 - \frac{2T}{n}\right) 2m'_t$. The constructions of the families guarantee that they have the same number of edges ($m_t = m'_t$) and different spectra.

■

The next corollary provides four non-isomorphic threshold graphs with same Laplacian energy.

Corollary 3.1. Consider the parameters t^b and t_1 of Theorems 2.1 and of Theorem 2.2, respectively. For all integer t with $1 \leq t \leq (n - T - 1)(T - 1)$, if $t \leq \min\{t^b, t_1\}$ then the graphs G_t in family $\mathcal{G}_{n,T}$, G_s in family $\mathcal{G}_{n,T'}$, F_t in family $\mathcal{F}_{n,T}$ and F_s in family $\mathcal{F}_{n,T'}$ are L -equienergetic, where $T' = n - T$ and $s = (n - T' - 1)(T' - 1) - (t - 1)$.

Proof: As in Theorem 3.1, manipulating algebraically some inequalities, we verify that if $t \leq t_1(T)$ then $s \geq t^\sharp(T')$. In addition, by the proof of Theorem 3.1, we have $s \geq t_2(T')$. Then Theorem 3.2 assures that the graphs G_s and F_s , of families $\mathcal{G}_{n,T'}$ and $\mathcal{F}_{n,T'}$, respectively, are L -equienergetic. But from Theorem 3.1, we know that $LE(F_s) = LE(F_t)$ (F_t graph in family $\mathcal{F}_{n,T}$). By using again Theorem 3.2, we obtain $LE(F_t) = LE(G_t)$ (G_t graph of $\mathcal{G}_{n,T}$), finally proving that $LE(G_s) = LE(F_s) = LE(F_t) = LE(G_t)$.

■

Example 1. Considering $n = 12$ and $T = 7$ we have $T' = 5$. Taking $t = 4$ we have $LE(G_4) = LE(F_4) = LE(F_{21}) = LE(G_{21}) = 44$, where $G_4 \in \mathcal{G}_{12,7}$, $F_4 \in \mathcal{F}_{12,7}$, $F_{21} \in \mathcal{F}_{12,5}$ and $G_{21} \in \mathcal{G}_{12,5}$.




Figure 4: G_4, F_4, G_{21}, F_{21} with $LE(G_4) = LE(F_4) = LE(F_{21}) = LE(G_{21}) = 44$.

References

- [CH77] Václav Chvátal and Peter L. Hammer, *Aggregation of inequalities in integer programming*, Studies in integer programming (Proc. Workshop, Bonn, 1975), Ann. of Discrete Math., vol. 1, North-Holland, Amsterdam, 1977, pp. 145–162. [MR 0479384](#)
- [FHT14] Eliseu Fritscher, Carlos Hoppen, and Vilmar Trevisan, *Unicyclic graphs with equal Laplacian energy*, Linear Multilinear Algebra **62** (2014), no. 2, 180–194. [MR 3175407](#)
- [GZ06] Ivan Gutman and Bo Zhou, *Laplacian energy of a graph*, Linear Algebra Appl. **414** (2006), no. 1, 29–37. [MR 2209232](#)
- [HZ77] Peter B. Henderson and Yechezkel Zalcstein, *A graph-theoretic characterization of the PV_{chunk} class of synchronizing primitives*, SIAM J. Comput. **6** (1977), no. 1, 88–108. [MR 0488948](#)
- [Mer94] Russell Merris, *Degree maximal graphs are Laplacian integral*, Linear Algebra Appl. **199** (1994), 381–389. [MR 1274427](#)
- [Ste09] Dragan Stevanović, *Large sets of noncospectral graphs with equal Laplacian energy*, MATCH Commun. Math. Comput. Chem. **61** (2009), no. 2, 463–470. [MR 2501828](#)

- [VDVJT13] Cybele T. M. Vinagre, Renata R. Del-Vecchio, Dagoberto A. R. Justo, and Vilmar Trevisan, *Maximum Laplacian energy among threshold graphs*, Linear Algebra Appl. **439** (2013), no. 5, 1479–1495. [MR 3067817](#)

R. R. Del-Vecchio
Instituto de Matemática e Estatística da UFF
Niterói, Rio de Janeiro, Brasil
renata@vm.uff.br

G. B. Pereira
Instituto de Matemática e Estatística da UFF
Niterói, Rio de Janeiro, Brasil
guilhermekbp@gmail.com

C. T. M. Vinagre
Instituto de Matemática e Estatística da UFF
Niterói, Rio de Janeiro, Brasil
cybl@vm.uff.br