

Spectral properties of KK_n^j graphs

Maria Freitas Renata Del-Vecchio  Nair Abreu * 

Abstract

Let G be a simple graph and $A = A(G)$, $L = L(G)$ and $Q = Q(G)$ the adjacency, the laplacian and the signless Laplacian matrices of G , respectively. For each of the associated matrices of G , $M = A, L$, or Q , we call M -spectrum of G the spectrum of the matrix M . In this work we present the KK_n^j graphs, obtained from two copies of the complete graph K_n by adding j edges, $1 \leq j \leq n$, between a vertex of one of the copies and j vertices of the other. We obtain M -spectral properties of this graph based on its clique number and its edge connectivity.

1 Introduction

Let $G = (V, E)$ be a simple graph on n vertices and $D(G) = \text{diag}(d_1, \dots, d_n)$ be the diagonal matrix of its vertex degrees. Let $A(G) = (a_{ij})$, where

$$a_{ij} = \begin{cases} 1, & \text{if } v_i \sim v_j \\ 0, & \text{otherwise.} \end{cases}$$

be the *adjacency* matrix of G . Let $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ be the *Laplacian* and the *signless Laplacian* matrices of G , respectively. For $M(G) = A(G)$, $L(G)$ or $Q(G)$, let $P_M(G, x)$ be the characteristic polynomial of $M(G)$ and $Sp_M(G)$ the spectrum of $M(G)$. A graph G is called M -integral when all eigenvalues of M are integer numbers. Since

2000 AMS Subject Classification. 68R10, 05C75, 05C85.

Key Words and Phrases: integral graph, characteristic polynomial of graph.

*Supported by CNPq.

1974, when Harary and Schwenk [6] posed the question *Which graphs have integral spectra?*, the search for A -integral graphs or L -integral graphs has been done. More recently, Q -integral graphs were introduced in the literature [1, 7, 8, 9, 4].

We recall that the clique number $\omega(G)$ of a graph G is the number of vertices in a maximum clique in G . The edge connectivity $\kappa'(G)$ of a graph G is the minimum number of edges whose deletion disconnects G .

In this work we obtain some spectral properties in a special class of graphs, based on these two important parameters.

2 Spectral properties of KK_n^j graphs

Definition 2.1. Let KK_n^j be the graph obtained from two copies of the complete graph K_n by adding j edges between one vertex of a copy of K_n and j vertices of the other copy, where j is such that $1 \leq j \leq n$.

Remark: This definition is a generalization of the graph KK_n^2 , introduced by Stevanović in [10], where it is analysed other spectral properties of this graph, as Laplacian energy.




Figure 1: KK_6^2

Remark: We note that $\omega(KK_n^j) = n$ and $\kappa'(KK_n^j) = j$ if $j \leq n - 1$ and $\kappa'(KK_n^j) = j - 1$ when $j = n$.

The next theorem gives explicitly the expression for the A -characteristic polynomial of KK_n^j :

Theorem 2.2. *If $1 \leq j \leq n$, $n \geq 3$, the A -characteristic polynomial of KK_n^j is $(x+1)^{(2n-4)}h(x)$, where $h(x) = x^4 + (4-2n)x^3 + (n^2-6n+6-j)x^2 + (2n^2-6n+2nj-j^2-3j+4)x + 1 + nj^2 - 2j^2 + n^2 - 2n - 2j + 3jn - jn^2$.*

Proof:

The adjacency matrix of KK_n^j can be represented in the form

$$A = \begin{bmatrix} \mathbb{J}_{n-1} - \mathbb{I}_{n-1} & \mathbf{1}_{n-1} & \mathbf{0}_{n-1,j} & \mathbf{0}_{n-1,n-j} \\ \mathbf{1}_{n-1}^T & 0 & \mathbf{1}_j^T & \mathbf{0}_{n-j}^T \\ \mathbf{0}_{j,n-1} & \mathbf{1}_j & \mathbb{J}_j - \mathbb{I}_j & \mathbb{J}_{j,n-j} \\ \mathbf{0}_{n-j,n-1} & \mathbf{0}_{n-j} & \mathbb{J}_{n-j,j} & \mathbb{J}_{n-j} - \mathbb{I}_{n-j} \end{bmatrix},$$

where \mathbb{J} is the matrix with all entries 1 and \mathbb{I} is the identity matrix.

We will now prove that -1 is an A -eigenvalue with multiplicity at least $2n-4$, exhibiting three kinds of eigenvectors corresponding to it.

We note that, for $\mathbf{u} \in \mathbb{R}^{n-1} \setminus \{0\}$ orthogonal to $\mathbf{1}_{n-1}$ and $\mathbf{v} = \begin{bmatrix} \mathbf{u} \\ \mathbf{0}_{n+1} \end{bmatrix} \in \mathbb{R}^{2n}$ we have that $A\mathbf{v} = -1\mathbf{v}$. Analogously, for $\mathbf{u} \in \mathbb{R}^{n-j} \setminus \{0\}$ orthogonal to $\mathbf{1}_{n-j}$, we have that $\mathbf{w} = \begin{bmatrix} \mathbf{0}_{n+j} \\ \mathbf{u} \end{bmatrix} \in \mathbb{R}^{2n}$ is such that $A\mathbf{w} = -1\mathbf{w}$.

For $\mathbf{u} \in \mathbb{R}^j \setminus \{0\}$ orthogonal to $\mathbf{1}_j$ and $\mathbf{v} = \begin{bmatrix} \mathbf{0}_n \\ \mathbf{u} \\ \mathbf{0}_{n-j} \end{bmatrix} \in \mathbb{R}^{2n}$ we have that $A\mathbf{v} = -1\mathbf{v}$. Again, -1 is an A -eigenvalue of KK_n^j with multiplicity at least $j-1$. Then, -1 is an A -eigenvalue of KK_n^j with multiplicity at least $2n-4$.

We note that the lines on each block of A have constant sum and we can consider now the matrix \overline{A} , where the entries are these sums:

$$\overline{A} = \begin{bmatrix} n-2 & 1 & 0 & 0 \\ n-1 & 0 & j & 0 \\ 0 & 1 & j-1 & n-j \\ 0 & 0 & j & n-j-1 \end{bmatrix}.$$

The characteristic polynomial of \overline{A} is $h(x) = x^4 + (4-2n)x^3 + (n^2-6n+6-j)x^2 + (2n^2-6n+2nj-j^2-3j+4)x + 1 + nj^2 - 2j^2 + n^2 - 2n - 2j + 3jn - jn^2$.

It is known that the eigenvalues of \overline{A} are eigenvalues of A , (see Theorem 2.1.3, page 5 in [3]) which completes the proof. ■

The conditions for A -integrality are:

Corollary 2.3. *The graph KK_n^j is A -integral if and only if the roots of $h(x)$ are integers.*

We now obtain the L -characteristic polynomial of KK_n^j :

Theorem 2.4. *If $1 \leq j \leq n$, $n \geq 3$, the L -characteristic polynomial of KK_n^j is $x(x-n)^{2n-j-2}(x-n-1)^{j-1}g(x)$, where $g(x) = x^2 - (n+1+j)x + 2j$.*

Proof: As in the previous theorem, the Laplacian matrix of KK_n^j can be represented in the form of a block matrix

$$L = \begin{bmatrix} (n)\mathbb{I}_{n-1} - \mathbb{J}_{n-1} & -\mathbf{1}_{n-1} & \mathbf{0}_{n-1,j} & \mathbf{0}_{n-1,n-j} \\ -\mathbf{1}_{n-1}^T & n-1+j & -\mathbf{1}_j^T & \mathbf{0}_{n-j}^T \\ \mathbf{0}_{j,n-1} & -\mathbf{1}_j & (n+1)\mathbb{I}_j - \mathbb{J}_j & -\mathbb{J}_{j,n-j} \\ \mathbf{0}_{n-j,n-1} & \mathbf{0}_{n-j} & -\mathbb{J}_{n-j,j} & (n)\mathbb{I}_{n-j} - \mathbb{J}_{n-j} \end{bmatrix}.$$

By an analogous reasoning, if $1 \leq j \leq n$, the L -eigenvalues of KK_n^j are 0, n with multiplicity $2n-j-2$, $n+1$ with multiplicity $j-1$, and $\frac{n+j+1 \pm \sqrt{(n+j+1)^2 - 8j}}{2}$

■.

In this case we obtain L -integrality conditions directly from the clique number and the edge connectivity of the graph.

Corollary 2.5. *The graph KK_n^j is L -integral if and only if $(n+1+j)^2 - 8j$ is a perfect square.*

Thus we can construct an infinite family of L -integral graphs:

Corollary 2.6. *For all $n \geq 3$, the graph KK_n^n is L -integral.*

In fact, in this case, the roots of $g(x)$ are 1 and $2n$.

Remarks on properties of L -spectrum:

Considering the family $\{KK_n^j : 1 \leq j \leq n\}$, we have that the graph KK_n^n behaves as an extremal graph according to the following aspects:

- The second smallest L -eigenvalue $a(G)$ of a graph G is called the algebraic connectivity of G . Fiedler, in [2] prove that for all G , $a(G) \leq k(G) \leq k'(G)$, where $k(G)$ denotes the vertex connectivity of G .

As $k(KK_n^j) = 1$ for any j , $1 \leq j \leq n$, $a(KK_n^j) \leq 1$. By Corollary 2.3, $a(KK_n^j)$ attains the maximum value, $a(KK_n^j) = 1$, when $j = n$ and so KK_n^n is an extremal graph with respect to the algebraic connectivity, for all n .

- It is well known that the greatest L -eigenvalue of a graph G , also called the L -index of the graph and denoted by $\mu_1(G)$, is always less than or equal to the size of the graph.

In our case, for any j , $1 \leq j \leq n$, $\mu_1(KK_n^j) \leq 2n$. Again, based on corollary 2.3, if $j = n$, the L -index attains its maximum value, $\mu_1(KK_n^j) = 2n$ and so KK_n^n is an extremal graph with respect to L -index, for all n .

Finally, we present the Q -characteristic polynomial of KK_n^j and some consequences of it, obtained in [5]

Theorem 2.7. *If $j \leq n \in \mathbb{N}$, $n \geq 3$, the Q -characteristic polynomial of KK_n^j is $(x - 2n + 2)(x - n + 1)^{j-1}(x - n + 2)^{2n-j-2}f(x)$, where $f(x) = x^2 - (3n + j + 3)x + 2(n^2 + n(j - 2) - 2j - 1)$.*

Consequently, the Q -eigenvalues of KK_n^j are:

- $2n - 2$,
- $n - 2$ with multiplicity $2n - j - 2$,
- $n - 1$ with multiplicity $j - 1$,
- $\frac{3n + j - 3 \pm \sqrt{(n - j - 1)^2 + 8j}}{2}$

Corollary 2.8. *The graph KK_n^j is Q -integral if and only if $(n - j - 1)^2 + 8j$ is a perfect square. Moreover, for $n, k, j \in \mathbb{N}$, if one of the conditions bellow is satisfied,*

1. $n = 3j$;
2. $n = 2j - 1$;
3. $n = 5k - 2$ and $j = 3k$;
4. $n = 3k + 6$ and $j = 2k + 6$;
5. $n = j = \frac{k(k+1)}{2}$.

then KK_n^j graph is Q -integral:

LQ -integral means L -integral and Q -integral. As an immediate consequence of these results and the characterization of the L -spectrum of the KK_n^j graph, we obtain an infinite family of LQ -integral graphs:

Corollary 2.9. *For all $k \in \mathbb{N}$, if $n = j = \frac{k(k+1)}{2}$, KK_n^j is an LQ -integral graph.*

3 Conclusion

For the three matrices considered, A , L or Q , we obtain the expressions of the characteristic polynomial based on $\omega(KK_n^j)$ and $\kappa'(KK_n^j)$. For $M = L$ or Q we have explicitly the M -spectrum in terms of these two parameters. Finally we built an infinite family of LQ -integral graphs. It remains open the question about the existence of A -integral graphs in the class considered.

References

- [1] D. Cvetković, P. Rowlinson, S. Simić, *Signless Laplacian of finite graphs*, Linear Algebra and its Applications **423** (2007), 155-171.
- [2] M. Fiedler: *Algebraic connectivity of graphs*. Czech. Math. J. 23 (98) (1973), 298-305.
- [3] M. A. A. de Freitas, Grafos integrais, laplacianos integrais e Q-integrais. (Portuguese), Ph. D. Thesis, UFRJ, Rio de Janeiro, 2009.
- [4] M.A.A. de Freitas, N.M.M. de Abreu, R.R. Del-Vecchio, S.Jurkiewicz, *Infinite families of Q-integral graphs*, Linear Algebra Appl. **432** (2010), 2353-2360.
- [5] M.A.A. de Freitas, N.M.M. de Abreu, R.R. Del-Vecchio, *Building infinite families of Q-integral graphs*, submitted.
- [6] F. Harary, A.J. Schwenk, *Which graphs have integral spectra?*, in Bari, R., Harary, F.(Eds), Graphs and Combinatorics, Springer, Berlin, (1974), 45-51.
- [7] S. Simić, Z. Stanić, *Q-integral graphs with edge-degrees at most five*, Discrete Math. **308** (2008), 4625-4634.
- [8] Z. Stanić, *There are exactly 172 connected Q-integral graphs up to 10 vertices*, Novi Sad J. Math. **37** (2) (2007), 193-205.
- [9] Z. Stanić, *Some results on Q-integral graphs*, Ars Combinatoria **90** (2009), 321-335.
- [10] D. Stevanović, I. Stanković, M. Milosević, *More on the relation between energy and Laplacian energy of graphs*, MATCH Commun. Math. Comput. Chem. 61 n. 2 (2009), 395-401.

Universidade Federal Fluminense
Instituto de Matemática e Estatística
24020-140 - Niterói, RJ - Brazil
Email: renata@vm.uff.br, rdelvecchio@ibest.com.br ,

Universidade Federal do Rio de Janeiro.
Centro de Tecnologia, Bloco F, Ilha do Fundão
22430-210 - Rio de Janeiro, RJ - Brazil.
E-Mail: nair@pep.ufrj.br